

BOSTER BIOLOGICAL TECHNOLOGY, LTD.

3942 B Valley Ave, Pleasanton, CA, 94566 Phone: 925-485-4527. 888-466-3604 Fax: 925-485-4560

www.bosterbio.com

Certificate of Analysis

Note: this is a sample COA. To get the COA for your lot #, please contact us at support@bosterbio.com

To whom it may concern,

This letter attests that Anti-FAK/PTK2 Antibody , catalog # PA1574, is manufactured by Boster Biological Technology., Ltd, through test and assay, we got the following data:

1. Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human FAK(1036-1052aa DVIDQARLKMLGQTRPH), identical to the related rat and mouse sequences.

2. Application Data

Applications Details

Western blot, 0.1-0.5 μ g/ml, Human, Rat, Mouse

3. Cross Reaction

No cross reactivity with other proteins

4. Image




Figure 1. Western blot analysis of FAK using anti-FAK antibody (PA1574).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human HEK293 whole cell lysates

Lane 2: human PC-3 whole cell lysates

Lane 3: human K562 whole cell lysates

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FAK antigen affinity purified polyclonal antibody (Catalog # PA1574) at 0.5 μ g/mL overnight at 4 $^{\circ}$ C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for FAK at approximately 119KD. The expected band size for FAK is at 119KD.




Figure 2. Western blot analysis of FAK using anti-FAK antibody (PA1574).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: rat heart tissue lysates

Lane 2: rat C6 whole cell lysates

Lane 3: mouse RAW246.7 whole cell lysates

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FAK antigen affinity purified polyclonal antibody (Catalog # PA1574) at 0.5 μ g/mL overnight at 4 $^{\circ}$ C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for FAK at approximately 119KD. The expected band size for FAK is at 119KD.

Approved by:

陈杨静

Ms. Yangjing Yue Chief technician

Quality Control System of ELISA

Date: May 13, 2016