

## Anti-TNF-Alpha (Q178) Antibody

Catalog Number: A00002

### About TNF

Anti-Glycogen Synthase 1 pS641 antibody is validated by IHC, Western Blot and ELISA. Human muscle glycogen synthase (GS) is responsible for the biosynthesis of glycogen from phosphorylated glucose units. Mammalian liver and muscle contain GS consisting of four subunits with a total molecular weight of 360,000. GS is subject to regulation through both allosteric and covalent modification and occurs in two forms: the phosphorylated inactive form, and the dephosphorylated active form. GS is inactivated by the serine/threonine kinase called glycogen synthase kinase-3 $\beta$  that mainly functions to phosphorylate muscle glycogen synthase. This antibody is specific for the phosphorylated form of GS at S641. Phosphorylation of GS at S641 has been associated with Antiphospholipid Antibody Syndrome.

### Overview


Product Name	Anti-TNF-Alpha (Q178) Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-TNF-Alpha (Q178) Antibody catalog # A00002. Tested in WB,IHC,IF applications. This antibody reacts with Human,Mouse,Rat.
Application	IF, IHC, WB
Clonality	Polyclonal PI9-17
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P01375

### Technical Details


Immunogen	Synthesized peptide derived from human p53 around the phosphorylation site of T81.
Predicted Reactive Species	Bovine, Chicken
Cross Reactivity	Weakly cross-reacts with dog p53.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-

	specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>WB: 1:500-1:2000</p> <p>IHC: 1:50-1:200</p> <p>IF: 1:200</p>

## Anti-TNF-Alpha (Q178) Antibody (A00002) Images



Immunohistochemistry (IHC) analyzes of TNF-alpha (Q178) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.



Western blot (WB) analysis of TNF-alpha (Q178) polyclonal antibody at 1:500 dilution Lane1:Hela cell lysate  
Lane2:NIH-3T3 cell lysate Lane3:Mouse heart tissue lysate  
Lane4:Rat brain tissue lysate

## 11 Publications Citing This Product

1. PubMed ID: 10.1016/j.lfs.2020.117292, Nifedipine inhibits oxidative stress and ameliorates osteoarthritis by activating the nuclear factor erythroid-2-related factor 2 pathway
2. PubMed ID: 10.1021/acsami.1c10309, Novel Multifunctional Silver Nanocomposite Serves as a Resistance-Reversal Agent to Synergistically Combat Carbapenem-Resistant *Acinetobacter baumannii*
3. PubMed ID: 10.1002/bem.22353, Effects of Ultrashort Wave Therapy on Inflammation and Macrophage Polarization after Acute Lung Injury in Rats

Visit [bosterbio.com/anti-tnf-alpha-q178-antibody-a00002-boster.html](https://bosterbio.com/anti-tnf-alpha-q178-antibody-a00002-boster.html) to see all 11 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TNF-Alpha (Q178) Antibody