

Anti-TNF alpha Antibody Picoband®

Catalog Number: A00002-2

About Tnf

TNFalpha (Tumor Necrosis Factor alpha) gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR2. And this cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. Moreover, this cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine.

Overview

Product Name	Anti-TNF alpha Antibody Picoband®
Reactive Species	Mouse
Description	Boster Bio Anti-TNF alpha Antibody Picoband® catalog # A00002-2. Tested in WB applications. This antibody reacts with Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , and 0.05 mg Na ₃ N. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P06804

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of mouse TNF alpha, different from the related human sequence by five amino acids, and from the related rat sequence by three amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.

Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Mouse

Anti-TNF alpha Antibody Picoband® (A00002-2) Images



Western blot analysis of TNF alpha using anti-TNF alpha antibody (A00002-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: mouse thymus tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TNF alpha antigen affinity purified polyclonal antibody (Catalog # A00002-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TNF alpha at approximately 25KD. The expected band size for TNF alpha is at 25KD.



Inflammation response were reduced by NLRC4 siRNA after ICH. a - g , j Western blot assay to detect NLRC4, pNLRC4, Pro-IL-1beta, IL-1beta, caspase-1, Pro-IL-18, TNF-alpha, and IL-6, (k , l) ELISA for IL-18 and IL-1beta (h , i) Immunostaining for MPO-positive cells ($\times 400$ and $\times 200$) at peri-hematoma area in sham, ICH, negative control siRNA, and NLRC4 siRNA groups at 72 h after ICH (six rats for each group). * P

40 Publications Citing This Product

1. PubMed ID: 10.1038/s12276-021-00702-y, The pivotal role of the NLRC4 inflammasome in neuroinflammation after intracerebral hemorrhage in rats
2. PubMed ID: 33607154, Liao L,Huang L,Wei X,Yin L,Wei X,Li T.Bioinformatic and biochemical studies of formononetin against liver injure.Life Sci.2021 Feb 16:119229.doi:10.1016/j.lfs.2021.119229.Epub ahead of print.PMID:33607154.
3. PubMed ID: 33607154, Liao L,Huang L,Wei X,Yin L,Wei X,Li T.Bioinformatic and biochemical studies of formononetin against liver injure.Life Sci.2021 Feb 16:119229.doi:10.1016/j.lfs.2021.119229.Epub ahead of print.PMID:33607154.

Visit [bosterbio.com/anti-tnf-alpha-picoband-trade-antibody-a00002-2-boster.html](https://www.bosterbio.com/anti-tnf-alpha-picoband-trade-antibody-a00002-2-boster.html) to see all 40 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TNF alpha Antibody

For Research Use Only. Not for use in diagnostic procedures.