

Anti-HIF-1 alpha/HIF1A Antibody Picoband®

Catalog Number: A00013-3

About HIF1A

HIF-1alpha (Hypoxia-inducible factor 1alpha, HIF1A) is a transcription factor that mediates cellular and systemic homeostatic responses to reduced O₂ availability in mammals, including angiogenesis, erythropoiesis and glycolysis. This gene was mapped to 14q21-q24. HIF-1alpha transactivate genes required for energy metabolism and tissue perfusion and is necessary for embryonic development and tumor explant growth. HIF-1alpha is over expressed during carcinogenesis, myocardial infarction and wound healing. It is crucial for the cellular response to hypoxia and is frequently over expressed in human cancers, resulting in the activation of genes essential for cell survival. HIF-1alpha regulates the survival and function in the inflammatory microenvironment directly. It is a transcription factor that plays a pivotal role in cellular adaptation to changes in oxygen availability.

Overview


Product Name	Anti-HIF-1 alpha/HIF1A Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-HIF-1 alpha/HIF1A Antibody Picoband® catalog # A00013-3. Tested in ELISA, Flow Cytometry, IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.01mg Na ₃ N.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q16665

Technical Details


Immunogen	E.coli-derived human HIF-1 alpha/HIF1A recombinant protein (Position: Q603-Q814).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Human, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5ug/ml, -


Anti-HIF-1 alpha/HIF1A Antibody Picoband® (A00013-3) Images



Western blot analysis of HIF-1 alpha/HIF1A using anti-HIF-1 alpha/HIF1A antibody (A00013-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions. Lane 1: human A431 whole cell lysates, Lane 2: human HEPG2 whole cell lysates, Lane 3: human U2OS whole cell lysates, Lane 4: rat PC-12 whole cell lysates, Lane 5: mouse thymus tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HIF-1 alpha/HIF1A antigen affinity purified polyclonal antibody (Catalog # A00013-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for HIF-1 alpha/HIF1A at approximately 93KD. The expected band size for HIF-1 alpha/HIF1A is at 93KD.




IHC analysis of HIF-1 alpha/HIF1A using anti-HIF-1 alpha/HIF1A antibody (A00013-3). HIF-1 alpha/HIF1A was detected in paraffin-embedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-HIF-1 alpha/HIF1A Antibody (A00013-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




IHC analysis of HIF-1 alpha/HIF1A using anti-HIF-1 alpha/HIF1A antibody (A00013-3). HIF-1 alpha/HIF1A was detected in paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-HIF-1 alpha/HIF1A Antibody (A00013-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.


IHC analysis of HIF-1 alpha/HIF1A using anti-HIF-1 alpha/HIF1A antibody (A00013-3). HIF-1 alpha/HIF1A was detected in paraffin-embedded section of mouse intestine



tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-HIF-1 alpha/HIF1A Antibody (A00013-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.



IHC analysis of HIF-1 alpha/HIF1A using anti-HIF-1 alpha/HIF1A antibody (A00013-3). HIF-1 alpha/HIF1A was detected in paraffin-embedded section of rat intestine tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-HIF-1 alpha/HIF1A Antibody (A00013-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.



Flow Cytometry analysis of A431 cells using anti-HIF-1 alpha/HIF1A antibody (A00013-3). Overlay histogram showing A431 cells stained with A00013-3 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-HIF-1 alpha/HIF1A Antibody (A00013-3, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

28 Publications Citing This Product

1. PubMed ID: 10.1590/1414-431X20142910, Putative role of ischemic postconditioning in a rat model of limb ischemia and reperfusion: involvement of hypoxia-inducible factor-1alpha expression
2. PubMed ID: 10.3748/wjg.v21.i7.2191, Semaphorin 4D and hypoxia-inducible factor-1alpha overexpression is related to prognosis in colorectal carcinoma
3. PubMed ID: 10.1186/s13046-016-0399-7, Hyperthermia induced HIF-1a expression of lung cancer through AKT and ERK signaling pathways

Visit bosterbio.com/anti-hif-1-alpha-hif1a-picoband-trade-antibody-a00013-3-boster.html to see all 28 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HIF-1 alpha/HIF1A Antibody

For Research Use Only. Not for use in diagnostic procedures.