

## Anti-IL10 Antibody

Catalog Number: A00021-3

### About IL10

Interleukin-10 (IL-10 or IL10), also known as human cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine. In humans IL-10 is encoded by the IL10 gene. It is capable of inhibiting synthesis of pro-inflammatory cytokines like IFN-gamma, IL-2, IL-3, TNFalpha and GM-CSF made by cells such as macrophages and regulatory T-cells. IL-10 also displays potent abilities to suppress the antigen presentation capacity of antigen presenting cells. Kim et al. (1992) showed that the mouse IL 10 gene contains 5 exons and spans about 5.2 kb of genomic DNA. Eskdale et al. (1997) mapped the IL10 gene to the junction between 1q31 and 1q32.

### Overview


Product Name	Anti-IL10 Antibody
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-IL10 Antibody catalog # A00021-3. Tested in ELISA, IHC applications. This antibody reacts with Mouse, Rat.
Application	ELISA, IHC
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P29456

### Technical Details


Immunogen	E.coli-derived rat IL10 recombinant protein (Position: K52-A170).
Recommended Detection Systems	Boster recommends HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Mouse, Rat

ELISA, 0.1-0.5ug/ml,


## Anti-IL10 Antibody (A00021-3) Images




Effects of Pseudoephedrine + emodin on the expression of TNF-alpha, IL-6, IL-1beta, iNOS, IL-10, Arg-1 in LPS-induced ALI rats. The contents of TNF-alpha ( A ), IL-6 ( B ), IL-1beta ( C ), iNOS ( D ), IL-10 ( E ), Arg-1 ( F ) in the serum and BALF were determined using ELISA. Data were expressed as mean  $\pm$  S.D. (n = 3). # p<0.05, ## p<0.01, ### p<0.001 vs. control group. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 vs. LPS alone group. + p<0.05, ++ p<0.01, +++ p<0.001 vs. combined treatment group (5 + 20 mg/kg) Index in PubMed under a CC BY license. PMID: 35123524



Effects of VIP on levels of inflammatory and anti-inflammatory cytokines in LPS + IFN-gamma or IL-4 + IL-13 induced AMs cells. Macrophages were pre-treated with VIP(10<sup>-6</sup>, 10<sup>-7</sup>, 10<sup>-8</sup> mol/L) for 24 h, followed by LPS (100 ng/mL) + IFN-gamma(20 ng/mL) or IL-4 (40 ng/mL) + IL-13(20 ng/mL) stimulation for 12 h. Cells were collected and the contents of TNF-alpha ( A ), IL-6 ( B ), IL-10 ( C ), Arg-1 ( D ) were determined using ELISA. (E-J) TNF-alpha, IL-6, iNOS, IL-10, Arg-1, Ym-1 mRNA expression was determined using Real-time PCR analysis. Data were expressed as mean  $\pm$  S.D. (n = 3). # p<0.05, ## p<0.01, ### p<0.001 vs. control group. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 vs. LPS alone group. +p<0.05, ++ p<0.01, +++ p<0.001 vs. IL-4 alone group Index in PubMed under a CC BY license. PMID: 35123524




VIP inhibited (M1) macrophage and activated (M2) macrophage in LPS + IFN-gamma-induced or IL-4 + IL-13-induced AMs. Effects of Pseudoephedrine + emodin on the expression of VIP in LPS-induced ALI rats. A - D , F , H , I (M1) macrophage Marker CD86, IL-6 and (M2) macrophage Marker CD206, IL-10 expression was determined using immunofluorescence. E , G CD86, CD206 mRNA expression was determined using Real-time PCR analysis. J The contents of VIP in the serum of ALI rats were determined using ELISA. Data are expressed as mean  $\pm$  S.D. (n = 3). ## p<0.01, ### p<0.001 vs. control group. \*p<0.05, \*\*p<0.01 vs. LPS alone group. +p<0.05, ++ p<0.01, +++ p<0.001 vs. IL-4 alone group Index in PubMed under a CC BY license. PMID: 35123524




Pseudoephedrine + emodin Inhibited MAPK in LPS-induced acute lung injury in rats. A - D Western blot analysis was performed to detect p-P38, p-ERK1/2 and p-JNK1/2 protein expression. E - I TNF-alpha, IL-6, IL-1beta, IL-10, Arg-1 mRNA expression was determined using Real-time PCR analysis. All data are expressed as mean  $\pm$  S.D. (n = 3). ## p<0.01, ### p<0.001 vs. control group. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 vs. LPS alone group. + p<0.05, ++ p<0.01, +++ p<0.001 vs. Combined treatment group (5 + 20 mg/kg) Index in PubMed under a CC BY license. PMID:


35123524




Effects of polysaccharide extract from XJEK on TNF-alpha, IL-1beta and IL-10 of HUVECs induced by Ang II. ( a ) TNF-alpha level in supernatants of HUVECs; ( b ) IL-1beta level in supernatants of HUVECs; ( c ) IL-10 level in supernatants of HUVECs. 1, blank control group; 2, Ang II (10 – 5 mol/L) group; 3, Ang II (10 – 5 mol/L) + AqE (0.15 mg/ml) group; 4, Ang II (10 – 5 mol/L) + AqE (0.3 mg/ml) group; 5, Ang II (10 – 5 mol/L) + AqE (0.6 mg/ml) group; 6, Ang II (10 – 5 mol/L) + AqE (1.2 mg/ml) group; 7, Ang II (10 – 5 mol/L) + XJEK (1.6 mg/ml) group. Data are expressed as mean ± SD, n = 6. \*\* P



Effects of polysaccharide extract from XJEK on TNF-alpha, IL-1beta and IL-10 in L -NAME-induced hypertensive mice. ( a ) TNF-alpha expression level in plasma. ( b ) IL-1beta expression level in plasma. ( c ) IL-10 expression level in plasma. ( d ) IL-1beta expression level in cardiac tissues. ( e ) TNF-alpha expression level in cardiac tissues. ( f ) IL-10 expression level in cardiac tissues. ( g ) Representative image of IL-1beta immunocytochemistry. ( h ) Representative image of TNF-alpha immunocytochemistry. ( i ) Representative image of IL-10 immunocytochemistry. 1, negative group; 2, control group; 3, model group; 4, L -NAME+AqE group; 5, L -NAME+XJEK group; 6, L -NAME+fisinopril group. Data are presented as the mean ± SD ( n = 10). \*\* P




SADS inhibited Raw264.7 and fibroblast-like synoviocyte inflammation in vitro. (A) The effect of SADS on the viability of Raw264.7. (B-D) RT-PCR analysis of TNF-alpha, IL-6 and IL-10 in Raw264.7 treated with SADS. (E) The effect of SADS on the viability of fibroblast-like synoviocytes. (F-H) RT-PCR analysis of TNF-alpha, IL-6 and IL-10 in fibroblast-like synoviocytes treated with SADS. (I) The phenotype of Raw264.7 was analyzed by flow cytometry. (J and K) Statistics of the proportion of M2 and M1 macrophages. (L and M) TNF-alpha protein expression level detection. ##P < 0.01 versus Ctrl; \*P < 0.01 versus LPS. n = 6. Index in PubMed under a CC BY license. PMID: 40688514




Pathological evaluation of IL-1RA–/– mice synovial tissue. HE staining showed synovial hyperplasia (A) and inflammation (B). (C) IHC staining of TNF-alpha in synovial tissue. (D) IHC staining of IL-10 in synovial tissue. (E and F) Levels of TNF-alpha and IL-10 in the blood of IL1RA–/– -deficient mice. ##P < 0.01 versus Ctrl; \*\*P < 0.01 versus Model. n = 6 mice for each group. Control is wild-type mice, and the model is IL1RA–/– mice. Index in PubMed under a CC BY license. PMID: 40688514

IHC analysis of IL10 using anti-IL10 antibody (A00021-3). IL10 was detected in paraffin-embedded section of mouse



spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-IL10 Antibody (A00021-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.



IHC analysis of IL10 using anti-IL10 antibody (A00021-3). IL10 was detected in paraffin-embedded section of rat spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-IL10 Antibody (A00021-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

## 7 Publications Citing This Product

1. PubMed ID: 10.1590/1414-431X20142910, Putative role of ischemic postconditioning in a rat model of limb ischemia and reperfusion: involvement of hypoxia-inducible factor-1alpha expression
2. PubMed ID: PMID:25337228, Human interleukin-10 gene inhibits acute rejection by triggering apoptosis in allograft vascular transplantation
3. PubMed ID: 10.1590/s0102-865020170050000007, Myocardial ischemic post-conditioning protects the lung against myocardial ischemia/reperfusion-induced damage by activating GSK-3beta

Visit [bosterbio.com/anti-il10-picoband-trade-antibody-a00021-3-boster.html](http://bosterbio.com/anti-il10-picoband-trade-antibody-a00021-3-boster.html) to see all 7 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IL10 Antibody

For Research Use Only. Not for use in diagnostic procedures.