

## Anti-Ubiquitin Antibody (N-term)

Catalog Number: A00060

### About UBC

This gene encodes ubiquitin, one of the most conserved proteins known. Ubiquitin is required for ATP-dependent, nonlysosomal intracellular protein degradation of abnormal proteins and normal proteins with a rapid turnover. Ubiquitin is covalently bound to proteins to be degraded, and presumably labels these proteins for degradation. Ubiquitin also binds to histone H2A in actively transcribed regions but does not cause histone H2A degradation, suggesting that ubiquitin is also involved in regulation of gene expression. This gene consists of three direct repeats of the ubiquitin coding sequence with no spacer sequence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. Aberrant form of this protein has been noticed in patients with Alzheimer's and Down syndrome.

### Overview


Product Name	Anti-Ubiquitin Antibody (N-term)
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Ubiquitin Antibody (N-term) (Catalog # A00060). Tested in IHC-P-Leica, WB application(s). This antibody reacts with Human, Mouse, Rat.
Application	WB, IHC-P-Leica
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P0CG48

### Technical Details


Immunogen	This Ubiquitin antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-32 amino acids from the N-terminal region of human Ubiquitin.
Predicted Reactive Species	Bovine, Chicken, Drosophila, Hamster, Horse, Monkey, Pig, Rabbit, Sheep, Xenopus
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid

Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:2000 IHC-P-Leica: 1:500


## Anti-Ubiquitin Antibody (N-term) (A00060) Images



All lanes : Anti-Ubiquitin Antibody (N-term) at 1:2000 dilution  
Lane 1: 293 whole cell lysate  
Lane 2: HeLa whole cell lysate  
Lane 3: HepG2 whole cell lysate  
Lane 4: A549 whole cell lysate  
Lane 5: NIH/3T3 whole cell lysate  
Lane 6: C2C12 whole cell lysate  
Lane 7: C6 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 10 kDa  
Blocking/Dilution buffer: 5% NFDm/TBST.



Immunohistochemical analysis of paraffin-embedded Human kidney tissue using A00060 performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody (1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded Human breast carcinoma tissue using A00060 performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody (1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Ubiquitin Antibody (N-term)

For Research Use Only. Not for use in diagnostic procedures.