

## Anti-CD11b/ITGAM Antibody Picoband®

Catalog Number: A00144-1

### About ITGAM

Integrin alpha M (ITGAM) is one protein subunit that forms the heterodimeric integrin alpha-M beta-2 (alphaMbeta2) molecule, also known as macrophage-1 antigen (Mac-1) or cluster of differentiation molecule 11B (CD11B). It is mapped to 16p11.2. ITGAM has a role in vascular repair after mechanical arterial injury. It is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles. What's more, ITGAM probably recognizes the R-G-D peptide in C3b, and it is also a receptor for fibrinogen, factor X and ICAM1.

### Overview

Product Name	Anti-CD11b/ITGAM Antibody Picoband®
Reactive Species	Human, Mouse
Description	Boster Bio Anti-CD11b/ITGAM Antibody Picoband® catalog # A00144-1. Tested in ELISA, Flow Cytometry, IHC, WB applications. This antibody reacts with Human, Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P11215


### Technical Details

Immunogen	E. coli-derived human CD11b recombinant protein (Position: N129-E336).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.


Suggested Dilutions

Western blot, 0.1-0.5ug/ml, Mouse  
Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human  
Flow Cytometry (Fixed), 1-3ug/1x10<sup>6</sup> cells, Human  
ELISA, 0.1-0.5ug/ml


## Anti-CD11b/ITGAM Antibody Picoband® (A00144-1) Images



Western blot analysis of CD11b using anti-CD11b antibody (A00144-1). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: mouse spleen tissue lysates, Lane 2: mouse RAW264.7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD11b antigen affinity purified polyclonal antibody (A00144-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for CD11b at approximately 179 kDa. The expected band size for CD11b is at 127 kDa.



IHC analysis of CD11b using anti-CD11b antibody (A00144-1). CD11b was detected in a paraffin-embedded section of human spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CD11b Antibody (A00144-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Flow Cytometry analysis of U937 cells using anti-CD11b antibody (A00144-1). Overlay histogram showing U937 cells stained with A00144-1 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-CD11b Antibody (A00144-1, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

## 5 Publications Citing This Product

1. PubMed ID: 25691929, Mu L, Jing C, Guo Z. Iran J Basic Med Sci. 2014 Nov;17(11):874-8. Expressions Of Cd11A, Cd11B, And Cd11C Integrin Proteins In Rats With Myocardial Hypertrophy.

2. PubMed ID: 28100749, Promoted interaction of C/EBP $\beta$ 1 with demethylated Cxcr3 gene promoter contributes to neuropathic pain in mice

3. PubMed ID: 20582224, Cloning, characterization, and expression analysis of goat (*Capra hircus*) phospholipid hydroperoxide glutathione peroxidase (PHGPx)

Visit [bosterbio.com/anti-cd11b-picoband-trade-antibody-a00144-1-boster.html](http://bosterbio.com/anti-cd11b-picoband-trade-antibody-a00144-1-boster.html) to see all 5 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD11b/ITGAM Antibody

For Research Use Only. Not for use in diagnostic procedures.