

Anti-NCAM1 Antibody Picoband™

Catalog Number: A00184

About NCAM1

NCAM is a membrane-bound glycoprotein that plays a role in cell-cell and cell-matrix adhesion through both its homophilic and heterophilic binding activity. The neural cell adhesion molecule appears on early embryonic cells and is important in the formation of cell collectives and their boundaries at sites of morphogenesis. Later in development it is found on various differentiated tissues and is a major CAM mediating adhesion among neurons and between neurons and muscle. NCAM gene is mapped to 11q23. The neural cell adhesion molecule (NCAM) can influence a number of diverse intercellular events, including junctional communication, the association of axons with pathways and targets, and signals that alter levels of neurotransmitter enzymes.

Overview

Product Name	Anti-NCAM1 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-NCAM1 Antibody Picoband™ catalog # A00184. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.01mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P13591

Technical Details

Immunogen	E.coli-derived human NCAM1 recombinant protein (Position: K152-Q690).
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.25ug/ml, Mouse, Rat</p> <p>Flow Cytometry(Fixed), 1-3 ug/1x10⁶ cells, Human</p> <p>Direct ELISA, 0.1-0.5ug/ml, Human</p>

Anti-NCAM1 Antibody Picoband™ (A00184) Images




Figure 1. Western blot analysis of NCAM1 using anti-NCAM1 antibody (A00184).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: rat brain tissue lysates,

Lane 2: rat C6 whole cell lysates,

Lane 3: mouse brain tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes.

Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NCAM1 antigen affinity purified polyclonal antibody (Catalog # A00184) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for NCAM1 at approximately 120-150KD. The expected band size for NCAM1 is at 95KD.




Figure 2. Flow Cytometry analysis of U937 cells using anti-NCAM1 antibody (A00184).

Overlay histogram showing U937 cells stained with A00184 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-NCAM1 Antibody (A00184, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

1 Publications Citing This Product

1. PubMed ID: 22460468, Assessment of the innate and adaptive immune system in proliferative vitreoretinopathy

Visit bosterbio.com/anti-ncam1-picoband-trade-antibody-a00184-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NCAM1 Antibody [™]