

Anti-CD86 Antibody

Catalog Number: A00220

About CD86

CD86, also known as B7-2, is a type I membrane protein that is a member of the immunoglobulin superfamily. Like the related protein CD80, this protein is expressed by antigen-presenting cells, and is the ligand for two proteins at the cell surface of T cells, CD28 and the cytotoxic T-lymphocyte-associated protein 4 (CTLA-4). Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell and induces T-cell proliferation and cytokine production. CTLA-4 binding negatively regulates T-cell activation and diminishes the immune response (1). Blocking the CTLA-4-CD80/CD86 interaction has been shown to enhance T-cell functions in acute lymphoblastic leukemia (ALL), suggesting that this pathway may be an attractive target for future cancer immunotherapy (2).

Overview


| | |
|----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name | Anti-CD86 Antibody |
| Reactive Species | Human |
| Description | Boster Bio Anti-CD86 Antibody (Catalog # A00220). Tested in ELISA, WB, IHC-P, IF applications. This antibody reacts with Human. |
| Application | ELISA, IF, IHC-P, WB |
| Clonality | Polyclonal |
| Formulation | CD86 Antibody is supplied in PBS containing 0.02% sodium azide. |
| Storage Instructions | CD86 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. Avoid repeated freeze-thaw cycles. Antibodies should not be exposed to prolonged high temperatures. |
| Host | Rabbit |
| Uniprot ID | P42082 |

Technical Details


| | |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunogen | CD86 antibody was raised against a peptide corresponding to 17 amino acids near the center of human CD86. The immunogen is located within amino acids 160 - 210 of CD86. |
| Predicted Reactive Species | Mouse, Rabbit |
| Cross Reactivity | At least five isoforms of CD86 are known to exist; this antibody will detect all five isoforms. |
| Isotype | IgG |
| Form | Liquid |
| Concentration | 1 mg/mL |

| | |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purification | CD86 Antibody is affinity chromatography purified via peptide column. |
| Suggested Dilutions | CD86 antibody can be used for Western blot at 1 - 2 ug/mL. Antibody can also be used for Immunohistochemistry at 2 ug/mL. For Immunofluorescence start at 20 ug/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher. |


Anti-CD86 Antibody (A00220) Images



Western blot analysis of CD86 in (A) Daudi and (B) Raji cell lysate with CD86 antibody at 1 ug/mL.



Immunohistochemistry of CD86 in human lymph node tissue with CD86 antibody at 2 ug/ml.



Immunofluorescence of CD86 in human lymph node tissue with CD86 antibody at 20 ug/ml. Red: CD86 Antibody (A00220) Blue: DAPI staining

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD86 Antibody

For Research Use Only. Not for use in diagnostic procedures.