

Anti-HDAC-1 (near C-terminus) Antibody

Catalog Number: A00256-2

About HDAC1

HDAC-1 antibody recognizes HDAC1 (also known as HD1, histone deacetylase 1, RPD3, RPD3L1) which belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis.

Overview

| | |
|----------------------|--|
| Product Name | Anti-HDAC-1 (near C-terminus) Antibody |
| Reactive Species | Human |
| Description | Boster Bio Anti-HDAC-1 (near C-terminus) Antibody (Catalog # A00256-2). Tested in ELISA, IF, IHC, WB applications. This antibody reacts with Human. |
| Application | ELISA, IF, IHC, WB |
| Clonality | Polyclonal |
| Formulation | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1% (w/v) Sodium Azide |
| Storage Instructions | Store vial at -20°C or below prior to opening. This vial contains a relatively low volume of reagent (25 uL). To minimize loss of volume dilute 1:10 by adding 225 uL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing. Expiration date is one (1) year from date of opening. (Ship on dry ice.) |
| Host | Rabbit |
| Uniprot ID | Q13547 |

Technical Details

| | |
|----------------------------|--|
| Immunogen | Anti-HDAC-1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a C-Terminal region near amino acids 450-482 of Human HDAC-1. |
| Predicted Reactive Species | Primate |
| Isotype | IgG |
| Form | Liquid (sterile filtered) |
| Concentration | 1.0 mg/mL by UV absorbance at 280 nm |

| | |
|---------------------|---|
| Purification | Anti-HDAC-1 antibody is directed against human HDAC-1 protein. HDAC-1 antibody was affinity purified from monospecific antiserum by immunoaffinity purification. A BLAST analysis was used to suggest reactivity with this protein from human, mouse, rat and chimpanzee sources based on 100% homology for the immunogen sequence. Cross-reactivity may occur with HDAC-1 from bovine (82% homology) and chicken (80% homology) sources. Cross-reactivity with HDAC-1 homologues from other sources has not been determined. |
| Suggested Dilutions | ELISA: 1:10,000 - 1:50,000 IHC: 1:200 - 1:1,000 IF Microscopy: 10ug / ml WB: 1:1,000 - 1:5,000 Anti-HDAC-1 Antibody has been tested for use in ELISA, immunohistochemistry, immunofluorescence, and western blot. Specific conditions for reactivity should be optimized by the end user. Specific nuclear staining is observed by IHC. Expect bands at 65 kDa in size corresponding to HDAC-1 by western blotting in the appropriate cell lysate or extract. |

1 Publications Citing This Product

1. PubMed ID: 25333250, Chen Xm, Xie Xb, Zhao Q, Wang F, Bai Y, Yin Jq, Jiang H, Xie Xi, Jia Q, Huang G. Mol Med Rep. 2015 Jan;11(1):105-12. Doi: 10.3892/Mmr.2014.2733. Epub 2014 Oct 21. Ampelopsin Induces Apoptosis By Regulating Multiple C-Myc/S-Phase Kinase-Associated ...

Visit bosterbio.com/anti-hdac-1-near-c-terminus-antibody-a00256-2-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HDAC-1 (near C-terminus) Antibody

For Research Use Only. Not for use in diagnostic procedures.