

Anti-TNFSF11 Antibody Picoband®

Catalog Number: A00363-2

About TNFSF11

Enables identical protein binding activity; receptor ligand activity; and tumor necrosis factor receptor superfamily binding activity. Involved in several processes, including intracellular signaling cassette; positive regulation of secretion by cell; and regulation of DNA-templated transcription. Acts upstream of or within several processes, including cellular response to cytokine stimulus; mammary gland development; and positive regulation of intracellular signal transduction. Predicted to be located in cytoplasm and plasma membrane. Predicted to be active in extracellular space. Is expressed in bone; genitourinary system; hemolymphoid system; incisor; and rib. Used to study autosomal recessive osteopetrosis 2 and osteoporosis. Human ortholog(s) of this gene implicated in autosomal recessive osteopetrosis 2 and osteoporosis. Orthologous to human TNFSF11 (TNF superfamily member 11).

Overview


Product Name	Anti-TNFSF11 Antibody Picoband®
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-TNFSF11 Antibody Picoband® catalog # A00363-2. Tested in WB, IHC, ELISA applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	O35235

Technical Details


Immunogen	E.coli-derived human TNFSF11 recombinant protein (Position: R72-D316).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Mouse, Rat

	Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Mouse, Rat ELISA, 0.1-0.5 ug/ml
--	--


Anti-TNFSF11 Antibody Picoband® (A00363-2) Images



Western blot analysis of TNFSF11 using anti-TNFSF11 antibody (A00363-2). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat thymus tissue lysates, Lane 2: rat C6 whole cell lysates, Lane 3: mouse spleen tissue lysates, Lane 4: mouse thymus tissue lysates, Lane 5: mouse colon tissue lysates, Lane 6: mouse Raw264.7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TNFSF11 antigen affinity purified polyclonal antibody (A00363-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for TNFSF11 at approximately 35 kDa. The expected band size for TNFSF11 is at 35 kDa.




IHC analysis of TNFSF11 using anti-TNFSF11 antibody (A00363-2). TNFSF11 was detected in a paraffin-embedded section of rat lymphaden tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TNFSF11 Antibody (A00363-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of TNFSF11 using anti-TNFSF11 antibody (A00363-2). TNFSF11 was detected in a paraffin-embedded section of rat lymphaden tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TNFSF11 Antibody (A00363-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

IHC analysis of TNFSF11 using anti-TNFSF11 antibody (A00363-2). TNFSF11 was detected in a paraffin-embedded section of mouse lymphaden tissue. Heat mediated antigen



retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TNFSF11 Antibody (A00363-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TNFSF11 Antibody

For Research Use Only. Not for use in diagnostic procedures.