

# Anti-Human TLR1 DyLight® 488 conjugated Antibody

Catalog Number: A00429-Dyl488

#### **About TLR1**

Toll-like receptor 1 (TLR1), also called TIL or CD281 is a member of the Toll-like receptor family (TLR) of pattern recognition receptors of the innate immune system. This gene is mapped to 4p14 by fluorescence in situ hybridization. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is ubiquitously expressed, and at higher levels than other TLR genes. Different length transcripts presumably resulting from use of alternative polyadenylation site, and/or from alternative splicing, have been noted for this gene.

#### Overview

Product Name	Anti-Human TLR1 DyLight® 488 conjugated Antibody
Reactive Species	Human
Description	Boster Bio Anti-Human TLR1 DyLight® 488 conjugated Antibody catalog # A00429-Dyl488. Tested in Flow Cytometry applications. This antibody reacts with Human.
Conjugate	DyLight®488
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na2HPO4, 0.02% NaN3.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q15399

#### **Technical Details**

Immunogen	E. coli-derived human TLR1 recombinant protein (Position: F350-D404).
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster provides a series of assays reacted with primary antibodies.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid



# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Concentration	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used: Flow Cytometry, 1-3ug/1x10 <sup>6</sup> cells



## Anti-Human TLR1 DyLight® 488 conjugated Antibody (A00429-Dyl488) Images



Boster Kit Box

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Human TLR1 DyLight® 488 conjugated Antibody