

## Anti-PPAR gamma/PPARG Antibody Picoband®

Catalog Number: A00449-2

### About PPARG

The peroxisome proliferator-activated receptors (PPARs) are a group of three nuclear receptor isoforms, PPAR gamma, PPAR alpha, and PPAR delta, encoded by different genes. PPARs are ligand-regulated transcription factors that control gene expression by binding to specific response elements (PPREs) within promoters. PPAR gamma is a transcription factor that has a pivotal role in adipocyte differentiation and expression of adipocyte-specific genes. The PPAR gamma1 and gamma2 isoforms result from alternative splicing and have ligand-dependent and -independent activation domains. PPAR gamma is a member of a family of nuclear receptors/ligand-dependent transcription factors, which bind to hormone response elements on target gene promoters. PPAR gamma is abundantly expressed in normal lung tissues, especially in endothelial cells, but that its expression is reduced or absent in the angiogenic plexiform lesions of pulmonary hypertensive lungs and in the vascular lesions of a rat model of severe pulmonary hypertension. And it is concluded that fluid shear stress decreases the expression of PPARgamma in endothelial cells and that loss of PPARgamma expression characterizes an abnormal, proliferating, apoptosis-resistant endothelial cell phenotype.

### Overview


Product Name	Anti-PPAR gamma/PPARG Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-PPAR gamma/PPARG Antibody Picoband® catalog # A00449-2. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P37231

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human PPAR gamma, identical to the related mouse and rat sequences.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.

Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human

## Anti-PPAR gamma/PPARG Antibody Picoband® (A00449-2) Images



Western blot analysis of PPAR gamma using anti-PPAR gamma antibody (A00449-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human K562 whole cell lysates, Lane 2: human MCF-7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PPAR gamma antigen affinity purified polyclonal antibody (Catalog # A00449-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PPAR gamma at approximately 58 kDa. The expected band size for PPAR gamma is at 58 kDa.

## 20 Publications Citing This Product

1. PubMed ID: 10.3892/etm.2011.208, Enhanced peroxisomal beta-oxidation metabolism in visceral adipose tissues of high-fat diet-fed obesity-resistant C57BL/6 mice
2. PubMed ID: 10.3389/fphys.2021.707344, Pioglitazone Ameliorates Renal Ischemia-Reperfusion Injury via Inhibition of NF-kappaB Activation and Inflammation in Rats
3. PubMed ID: 10.1186/s12883-015-0287-3, Rosiglitazone infusion therapy following minimally invasive surgery for intracerebral hemorrhage evacuation decreases matrix metalloproteinase-9 and blood-brain barrier disruption in rabbits

Visit [bosterbio.com/anti-ppar-gamma-picoband-trade-antibody-a00449-2-boster.html](http://bosterbio.com/anti-ppar-gamma-picoband-trade-antibody-a00449-2-boster.html) to see all 20 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PPAR gamma/PPARG Antibody

For Research Use Only. Not for use in diagnostic procedures.