

Anti-TRAIL/Tnfsf10 Antibody Picoband™

Catalog Number: A00466

About Tnfsf10

TNF-related apoptosis-inducing ligand (TRAIL), is a protein functioning as a ligand that induces the process of cell death called apoptosis. TRAIL has also been designated CD253 (cluster of differentiation 253). It is a member of Tumor Necrosis Factor Ligand Superfamily. In humans, the gene that encodes for TRAIL is located at chromosome 3q26. TRAIL has got 281 amino acid proteins. TRAIL mRNA is expressed at significant levels in most normal tissues. TRAIL binds to the death receptors DR4 (TRAIL-RI) and DR5 (TRAIL-RII) and the receptors DcR1 and DcR2, leading to activation of specific kinases and transcription of genes.

Overview

Product Name	Anti-TRAIL/Tnfsf10 Antibody Picoband™
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-TRAIL/Tnfsf10 Antibody Picoband™ catalog # A00466. Tested in WB applications. This antibody reacts with Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P50592

Technical Details

Immunogen	E.coli-derived mouse TRAIL recombinant protein (Position: Y39-N291). Mouse TRAIL shares 66.3% amino acid (aa) sequence identity with human TRAIL.
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Mouse, Rat</p>

Anti-TRAIL/Tnfsf10 Antibody Picoband™ (A00466) Images




Figure 1. Western blot analysis of TRAIL using anti-TRAIL antibody (A00466).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: HEPA1-6 whole Cell lysates,

Lane 2: NIH3T3 whole Cell lysates,

Lane 3: PC12 whole Cell lysates,

Lane 4: NRK whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes.

Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TRAIL antigen affinity purified polyclonal antibody (Catalog # A00466) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TRAIL at approximately 20KD. The expected band size for TRAIL is at 35KD.

1 Publications Citing This Product

1. PubMed ID: 28377479, AAV-iRFP labelling of human mesenchymal stem cells for near-infrared fluorescence imaging

Visit bosterbio.com/anti-trail-picoband-trade-antibody-a00466-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TRAIL/Tnfsf10 Antibody™