

Anti-MERTK Antibody

Catalog Number: A00489-3

About MERTK

MERTK is a member of the MER/AXL/TYRO3 receptor kinase family and a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type (immunoglobulin-like) domains, and one tyrosine kinase domain. Mutations in MERTK gene have been associated with disruption of the retinal pigment epithelium (RPE) phagocytosis pathway and onset of autosomal recessive retinitis pigmentosa (RP).

Overview

Product Name	Anti-MERTK Antibody
Reactive Species	Human
Description	Boster Bio Anti-MERTK Antibody (Catalog # A00489-3). Tested in WB, IF, Flow Cytometry, IHC-P application(s). This antibody reacts with Human.
Application	Flow Cytometry, IF, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q12866


Technical Details

Immunogen	This MERTK antibody is generated from rabbits immunized with a his tag recombinant protein of human MERTK.
Predicted Reactive Species	Mouse
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Purification	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.


Suggested Dilutions

IF: 1:10-1:50
WB: 1:1000
IHC-P: 1:50
FC: 1:10-1:50


Anti-MERTK Antibody (A00489-3) Images




Fluorescent image of U251 cell stained with MERTK Antibody. U251 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with MERTK primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). MERTK immunoreactivity is localized to Vesicles significantly.



MERTK Antibody western blot analysis in Jurkat cell line lysates (35 µg/lane). This demonstrates the MERTK antibody detected the MERTK protein (arrowhead).



Formalin-fixed and paraffin-embedded human lymphoma reacted with MERTK Antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of Jurkat cells using MERTK Antibody (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MERTK Antibody

For Research Use Only. Not for use in diagnostic procedures.