

## Anti-DLK1 Antibody (C-term)

Catalog Number: A00513-3

### About DLK1

May have a role in neuroendocrine differentiation.

### Overview


Product Name	Anti-DLK1 Antibody (C-term)
Reactive Species	Human, Mouse
Description	Boster Bio Anti-DLK1 Antibody (C-term) (Catalog # A00513-3). Tested in WB, Flow Cytometry, IHC-P application(s). This antibody reacts with Human, Mouse.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P80370

### Technical Details


Immunogen	This DLK1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 368-402 amino acids from the C-terminal region of human DLK1.
Predicted Reactive Species	Bovine, Rat, Xenopus
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:1000 IHC-P: 1:25 FC: 1:25




## Anti-DLK1 Antibody (C-term) (A00513-3) Images



All lanes : Anti-DLK1 Antibody (C-term) at 1:1000 dilution  
Lane 1: human placenta lysates  
Lane 2: mouse skeletal muscle lysates  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution  
Predicted band size : 41 kDa  
Blocking/Dilution buffer: 5% NFD/MTBST.



Immunohistochemical analysis of paraffin-embedded H. placenta section using DLK1 Antibody (C-term). A00513-3 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Flow cytometric analysis of HepG2 cells using DLK1 Antibody (C-term)(green, Cat#A00513-3) compared to an isotype control of mouse IgG2b(blue). A00513-3 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-DLK1 Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.