

Anti-GSTM1 Antibody (C-term)

Catalog Number: A00569

About GSTM1

GSTM1 is a glutathione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs.

Overview

Product Name	Anti-GSTM1 Antibody (C-term)
Reactive Species	Human, Mouse
Description	Boster Bio Anti-GSTM1 Antibody (C-term) (Catalog # A00569). Tested in WB, Flow Cytometry, IHC-P application(s). This antibody reacts with Human, Mouse.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P09488


Technical Details

Immunogen	This GSTM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 184-211 amino acids from the C-terminal region of human GSTM1.
Predicted Reactive Species	C. elegans
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.


Suggested Dilutions

WB: 1:1000
IHC-P: 1:10-1:50
FC: 1:10-1:50


Anti-GSTM1 Antibody (C-term) (A00569) Images



Western blot analysis of lysates from Jurkat, U251 cell line and human liver tissue (from left to right), using GSTM1 Antibody (C-term) . A00569 was diluted at 1:1000 at each lane. A goat anti-rabbit (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with GSTM1 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



GSTM1 Antibody (C-term) (Cat. #A00569) flow cytometric analysis of MDA-MB231 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-GSTM1 Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.