

## Anti-Lactoferrin/LTF Antibody Picoband®

Catalog Number: A00633-1

### About LTF

Lactoferrin (LF), also known as lactotransferrin (LTF), is a multifunctional protein of the transferrin family. The protein is a major iron-binding protein in milk and body secretions with an antimicrobial activity, making it an important component of the non-specific immune system. The protein demonstrates a broad spectrum of properties, including regulation of iron homeostasis, host defense against a broad range of microbial infections, anti-inflammatory activity, regulation of cellular growth and differentiation and protection against cancer development and metastasis. Antimicrobial, antiviral, antifungal and antiparasitic activity has been found for this protein and its peptides.

### Overview


Product Name	Anti-Lactoferrin/LTF Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Lactoferrin/LTF Antibody Picoband® catalog # A00633-1. Tested in ELISA, IF, IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, IF, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P02788

### Technical Details


Immunogen	E. coli-derived human Lactoferrin recombinant protein (Position: D529-K710).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human Immunofluorescence, 5ug/ml, Human ELISA, 0.1-0.5ug/ml, -


## Anti-Lactoferrin/LTF Antibody Picoband® (A00633-1) Images



Western blot analysis of LTF using anti-LTF antibody (A00633-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human hepatocellular carcinoma paracancerous tissue (HCCP) lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-LTF antigen affinity purified polyclonal antibody (A00633-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for LTF at approximately 85 kDa. The expected band size for LTF is at 78 kDa.




IHC analysis of LTF using anti-LTF antibody (A00633-1). LTF was detected in a paraffin-embedded section of human prostatic cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-LTF Antibody (A00633-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




LTF, a ferroptosis-related gene, is identified in aging cochleae. (A) The volcano diagram of shows 43385 DEGs between the old group ( ~864316) and young group ( ~864308). (B) The heatmap shows 28 statistically significant DEGs. (C) Functional analysis of the statistically significant DEGs. (D) LTF, the hub gene, is obtained from the intersection of Up-Gens, Down-Gens (based on the statistically significant DEGs), and FRGs. (E) The PPI network shows that 12 proteins are interacting with LTF. Index in PubMed under a CC BY license. PMID: 38282692


TF-miRNA-mRNA network in cochlear ferroptosis. (A) The volcano diagram of shows 7512 DEMs between the old group ( ~1095954) and young group ( ~1095948). (B) The heatmap shows 12 statistically significant DEMs. (C-F) TFs predicted by mmu-mir-130b (C) , mmu-mir-205 (D) , hsa-mir-130b (E) , and hsa-mir-205 (F) . (G) CEBPA and CEBPB were predicted by mir-130b. (H) STAT3 was predicted by mir-205. (I,J) The top 20 TFs predicted by LTF in mice (I) and




humans (J) , CEBPA (Red box) was identified through the intersection of (I,J) , and TFs predicted by miRNAs in (G,H) . (K) The regulatory network was constructed. Index in PubMed under a CC BY license. PMID: 38282692




Ferroptosis in aging HEI-OC1 cells. (A) SA-beta-gal staining in HEI-OC1 cells treated with different concentrations of D-gal for 48 h. (B) Quantification of SA-beta-positive cells in (A) . Compared with the control group, the percentage of positive cells is observed to have a statistical difference at 20 mg/mL D-gal ( p



Ferroptosis in aging cochlear explants. (A) SA-beta-gal staining in the basement membrane treated with different concentrations of D-gal for 48 h. (B) Quantification of SA-beta-positive cells in (A) . Compared with the control group, the percentage of positive cells is observed to have a statistical difference at 30 mg/mL D-gal ( p < 0.001; N = 3). (C-E) Compared with the control group, the expression of Fe 2+ (C) and MDA (D) is increased in aging cochlear explants with 30 mg/mL D-gal ( p




Ferroptosis in aging cochleae. (A) Compared with young mice (2mo), the distinctly increased ABR thresholds are shown in old mice (10mo) at all frequencies ( p



IF analysis of LTF using anti-LTF antibody (A00633-1). LTF was detected in a paraffin-embedded section of human prostatic cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-LTF Antibody (A00633-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Western blot analysis of Lactoferrin/LTF using anti-Lactoferrin/LTF Antibody (A00633-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing



conditions. Lane 1: Total LTF from mouse neutrophil cell lysates. Lane 2: degranulated LTF. Ratio 1:20. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% BSA for 1.5 hour at RT. The membrane was incubated with rabbit anti-Lactoferrin/LTF antigen affinity purified polyclonal antibody (A00633-1) at 1:2000 1h overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:3000 dilution at RT for 1 hour. The signal is developed using HRP. A specific band was detected for Lactoferrin/LTF at approximately 77 kDa. The expected band size for Lactoferrin/LTF is at 77 kDa.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-Lactoferrin/LTF Antibody

For Research Use Only. Not for use in diagnostic procedures.