

Anti-Cxcr5 Antibody

Catalog Number: A00663-1

About Cxcr5

CXCR5(Chemokine CXC Motif Receptor 5), also known as BLR1, is a G protein-coupled seven transmembrane receptor for chemokine CXCL13(also known as BLC) and belongs to the CXC chemokine receptor family. In humans, the CXC-R5 protein is encoded by the CXCR5 gene. The gene plays an essential role in B cell migration. Reif et al.(2002) concluded that their findings defined the mechanism of B-cell relocalization in response to antigen, and established that cell position in vivo can be determined by the balance of responsiveness to chemoattractants made in separate but adjacent zones. Chan et al.(2003) investigated the expression of chemokines and chemokine receptors in eyes with primary intraocular B-cell lymphoma(PIOL).

Overview

Product Name	Anti-Cxcr5 Antibody
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-Cxcr5 Antibody catalog # A00663-1. Tested in ELISA, IHC applications. This antibody reacts with Mouse, Rat.
Application	ELISA, IHC
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q04683


Technical Details

Immunogen	E.coli-derived mouse Cxcr5 recombinant protein (Position: M1-F374).
Recommended Detection Systems	Boster recommends HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.


Suggested Dilutions

Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Mouse, Rat
ELISA, 0.1-0.5 ug/ml, -

Anti-Cxcr5 Antibody (A00663-1) Images



IHC analysis of TFEB using anti-TFEB antibody (A00663-1). TFEB was detected in a paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TFEB Antibody (A00663-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of TFEB using anti-TFEB antibody (A00663-1). TFEB was detected in a paraffin-embedded section of rat lymph nodes tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TFEB Antibody (A00663-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cxcr5 Antibody

For Research Use Only. Not for use in diagnostic procedures.