

# **Anti-FGF8 Antibody Picoband™**

Catalog Number: A00841

### **About FGF8**

Fibroblast growth factor 8 (androgen-induced), also known as FGF8 or AIGF, is a human gene which maps to 10q24. The protein encoded by this gene is secreted proteins that interact with FGF tyrosine kinase receptors to mediate growth and development. This protein is known to be a factor that supports androgen and anchorage independent growth of mammary tumor cells. Overexpression of this gene has been shown to increase tumor growth and angiogensis. The temporal and spatial patterns of this gene expression suggest that FGF8 is involved in gastrulation, regionalization of the brain, and organogenesis of the limb and face as an embryonic epithelial factor. The adult expression of FGF8 is restricted to gonads, including testes and ovaries. FGF8 stimulated growth of human prostate carcinoma cells and mouse fibroblasts and mammary carcinoma cells in a dose-dependent manner. It also may play an important role in growth and patterning of limbs, face, and central nervous system. FGF8 is expressed in increased levels in breast cancer and in lactating human breast; it was also detected in human milk. A survey of other normal tissues showed that FGF8 is expressed in the proliferative cells of the skin and epithelial cells in colon, ovary, fallopian tube, and uterus.

#### Overview

Product Name	Anti-FGF8 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-FGF8 Antibody Picoband™ catalog # A00841. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	FGF8: P55075

#### **Technical Details**

Immunogen	E. coli-derived human FGF8 recombinant protein (Position: Q23-R233).
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG





888-466-3604 | support@bosterbio.com | www.bosterbio.com

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Western blot, 0.1-0.5ug/ml  Direct ELISA, 0.1-0.5ug/ml



## Anti-FGF8 Antibody Picoband™ (A00841) Images




Figure 1. Western blot analysis of FGF8 using anti-FGF8 antibody (A00841).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: rat ovary tissue lysates,

Lane 2: human Hela whole cell lysates,

Lane 3: human placenta tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FGF8 antigen affinity purified polyclonal antibody (Catalog # A00841) at 0.5 ug/mL overnight at 4 then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for FGF8 at approximately 26KD. The expected band size for FGF8 is at 26KD.

### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-FGF8 Antibody ™