

Anti-CD137/Tnfrsf9 Antibody Picoband®

Catalog Number: A00867-2

About Tnfrsf9

CD137 is a member of the tumor necrosis factor (TNF) receptor family. This receptor contributes to the clonal expansion, survival, and development of T cells. It can also induce proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 co-stimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-kappaB.

Overview

Product Name	Anti-CD137/Tnfrsf9 Antibody Picoband®
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-CD137/Tnfrsf9 Antibody Picoband® catalog # A00867-2. Tested in ELISA, WB applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB, ELISA (Cap)
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P20334


Technical Details

Immunogen	E. coli-derived mouse CD137 recombinant protein (Position: V24-L187).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.


Suggested Dilutions

Western blot, 0.1-0.5ug/ml
ELISA (Cap), 1-5ug/ml

Anti-CD137/Tnfrsf9 Antibody Picoband® (A00867-2) Images



Western blot analysis of CD137 using anti-CD137 antibody (A00867-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: recombinant mouse CD137 protein 1ng. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD137 antigen affinity purified polyclonal antibody (Catalog # A00867-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CD137 at approximately 65KD. The expected band size for CD137 is at 44KD.



Sandwich ELISA - Recombinant mouse CD137/Tnfrsf9 protein standard curve. Use in combination with reagents from Mouse CD137/Tnfrsf9 ELISA Kit EZ-Set (DIY Antibody Pairs) (EZ1143).

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD137/Tnfrsf9 Antibody

For Research Use Only. Not for use in diagnostic procedures.