

Anti-Epoxide hydrolase/EPHX1 Antibody Picoband®

Catalog Number: A00899-1

About EPHX1

Epoxyde hydrolase 1 is an enzyme encoded by the EPHX1 gene in humans. Epoxyde hydrolase is a critical biotransformation enzyme that converts epoxydes from the degradation of aromatic compounds to trans-dihydrodiols which can be conjugated and excreted from the body. Epoxyde hydrolase functions in both the activation and detoxification of epoxydes. Mutations in this gene cause preeclampsia, epoxyde hydrolase deficiency or increased epoxyde hydrolase activity. Alternatively spliced transcript variants encoding the same protein have been found for this gene.

Overview


Product Name	Anti-Epoxyde hydrolase/EPHX1 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Epoxyde hydrolase/EPHX1 Antibody Picoband® catalog # A00899-1. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P07099

Technical Details


Immunogen	E.coli-derived human Epoxyde hydrolase/EPHX1 recombinant protein (Position: A42-Q455).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3 ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5 ug/ml, -


Anti-Epoxyde hydrolase/EPHX1 Antibody Picoband® (A00899-1) Images



Western blot analysis of Epoxyde hydrolase/EPHX1 using anti-Epoxyde hydrolase/EPHX1 antibody (A00899-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human HCCT tissue lysates, Lane 3: human HCCP tissue lysates, Lane 4: rat liver tissue lysates, Lane 5: rat RH35 whole cell lysates, Lane 6: mouse liver tissue lysates, Lane 7: mouse HEP1-6 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Epoxyde hydrolase/EPHX1 antigen affinity purified polyclonal antibody (Catalog # A00899-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Epoxyde hydrolase/EPHX1 at approximately 53 kDa. The expected band size for Epoxyde hydrolase/EPHX1 is at 53 kDa.




Flow Cytometry analysis of RT4 cells using anti-Epoxyde hydrolase/EPHX1 antibody (A00899-1). Overlay histogram showing RT4 cells stained with A00899-1 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Epoxyde hydrolase/EPHX1 Antibody (A00899-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.



Flow Cytometry analysis of A549 cells using anti-Epoxyde hydrolase/EPHX1 antibody (A00899-1). Overlay histogram showing A549 cells stained with A00899-1 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Epoxyde hydrolase/EPHX1 Antibody (A00899-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody

(Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.



Flow Cytometry analysis of SiHa cells using anti-Epoxyde hydrolase/EPHX1 antibody (A00899-1). Overlay histogram showing SiHa cells stained with A00899-1 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Epoxyde hydrolase/EPHX1 Antibody (A00899-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Epoxyde hydrolase/EPHX1 Antibody

For Research Use Only. Not for use in diagnostic procedures.