

## Anti-HDAC-4 (internal) Antibody

Catalog Number: A00971-2

### About HDAC4

HDAC4 is a 140 kD class II histone deacetylase similar to the yeast HDA1 protein. This nuclear protein operates in concert with histone acetyltransferases to control core histone acetylation in highly conserved lysine residues. HDAC activity has been associated with transcriptional repression and nucleosomal condensation. HDAC4 uses active nuclear export to shuttle between the nucleus and the cytoplasm. HDAC4 is modified by phosphorylation by CaM kinase I or IV. HDAC4 has been reported to interact with 14-3-3 in the cytoplasm and MEF2A, HDAC3, NcoR and SMRT in the nucleus. Anti-HDAC4 antibodies are ideal for researchers interested in Breast Cancer, Cancer, Cell Cycle and Replication, Chromatin Research, Epigenetics, and Histone Deacetylases research.

### Overview


Product Name	Anti-HDAC-4 (internal) Antibody
Reactive Species	Human, Mouse, Rat, Primate
Description	Boster Bio Anti-HDAC-4 (internal) Antibody (Catalog # A00971-2). Tested in WB applications. This antibody reacts with Human, Mouse, Rat, Primate.
Application	ELISA, IF, IHC, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 50% (v/v) Glycerol
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is six (6) months from date of opening. (Ship on dry ice.)
Host	Rabbit
Uniprot ID	P56524

### Technical Details

Immunogen	HDAC4 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to the internal region surrounding 600-650aa of human HDAC4.
Predicted Reactive Species	African Green Monkey, Chimpanzee, Zebrafish
Isotype	IgG
Form	Liquid (sterile filtered)

Concentration	1 mg/mL by UV absorbance at 280 nm
Purification	Anti-HDAC4 was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody is specific towards HDAC4. A BLAST analysis was used to suggest cross-reactivity with Human, Mouse, Rat, and Primate based on 100% sequence homology. Cross-reactivity with HDAC4 from other sources has not been determined.
Suggested Dilutions	ELISA: 1:20,000 - 1:60,000 IHC: 1 mg/mL IF Microscopy: 1:100-1:500 WB: 1 mg/mL Anti-HDAC4 antibody has been tested in Western Blot and is useful for ELISA and immunohistochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~140 kDa corresponding to the appropriate cell lysate or extract.

## Anti-HDAC-4 (internal) Antibody (A00971-2) Images



Western Blot of Rabbit anti-HDAC4 antibody. Lane 1: Mouse brain homogenate. Lane 2: Mouse brain homogenate blocked with peptide. Load: 30  $\mu$ g per lane. Primary antibody: HDAC4 antibody at 1:400 for overnight at 4°C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 140 kDa, 98 kDa, and 55 kDa for HDAC4. Other band(s): HDAC4 splice variants and isoforms.

## 1 Publications Citing This Product

1. PubMed ID: 26473953, Arsenic trioxide reduces global histone H4 acetylation at lysine 16 through direct binding to histone acetyltransferase hMOF in human cells

Visit [bosterbio.com/anti-hdac-4-internal-antibody-a00971-2-boster.html](http://bosterbio.com/anti-hdac-4-internal-antibody-a00971-2-boster.html) to see all 1 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HDAC-4 (internal) Antibody

For Research Use Only. Not for use in diagnostic procedures.