

Anti-Myelin Protein Zero/MPZ Antibody Picoband™

Catalog Number: A00997-1

About MPZ

Myelin protein zero (P0, MPZ) is a glycoprotein which in humans is encoded by the MPZ gene. This gene is specifically expressed in Schwann cells of the peripheral nervous system and encodes a type I transmembrane glycoprotein that is a major structural protein of the peripheral myelin sheath. The encoded protein contains a large hydrophobic extracellular domain and a smaller basic intracellular domain, which are essential for the formation and stabilization of the multilamellar structure of the compact myelin. Mutations in this gene are associated with autosomal dominant form of Charcot-Marie-Tooth disease type 1 (CMT1B) and other polyneuropathies, such as Dejerine-Sottas syndrome (DSS) and congenital hypomyelinating neuropathy (CHN). A recent study showed that two isoforms are produced from the same mRNA by use of alternative in-frame translation termination codons via a stop codon readthrough mechanism.

Overview

Product Name	Anti-Myelin Protein Zero/MPZ Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Myelin Protein Zero/MPZ Antibody Picoband™ catalog # A00997-1. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P25189

Technical Details

Immunogen	E. coli-derived human Myelin Protein Zero recombinant protein (Position: I30-R153).
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized





Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml Direct ELISA, 0.1-0.5ug/ml



Anti-Myelin Protein Zero/MPZ Antibody Picoband™ (A00997-1) Images



Figure 1. Western blot analysis of Myelin Protein Zero using anti-Myelin Protein Zero antibody (A00997-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human U-87MG whole cell lysates.

After Electrophoresis, proteins were transferred to a
Nitrocellulose membrane at 150mA for 50-90 minutes.

Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5
hour at RT. The membrane was incubated with rabbit antiMyelin Protein Zero antigen affinity purified polyclonal
antibody (Catalog # A00997-1) at 0.5 ug/mL overnight at
4°C, then washed with TBS-0.1%Tween 3 times with 5
minutes each and probed with a goat anti-rabbit IgG-HRP
secondary antibody at a dilution of 1:10000 for 1.5 hour at
RT. The signal is developed using an Enhanced
Chemiluminescent detection (ECL) kit (Catalog # EK1002)
with Tanon 5200 system. A specific band was detected for
Myelin Protein Zero at approximately 28KD. The expected
band size for Myelin Protein Zero is at 28KD.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Myelin Protein Zero/MPZ Antibody ™