

# Anti-KCNH1 Antibody Picoband™

Catalog Number: A01036-2

#### **About KCNH1**

Potassium voltage-gated channel subfamily H member 1 is a protein that in humans is encoded by the KCNH1 gene. Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, subfamily H. This member is a pore-forming (alpha) subunit of a voltage-gated non-inactivating delayed rectifier potassium channel. It is activated at the onset of myoblast differentiation. The gene is highly expressed in brain and in myoblasts. Overexpression of the gene may confer a growth advantage to cancer cells and favor tumor cell proliferation. Alternative splicing of this gene results in two transcript variants encoding distinct isoforms.

### Overview

| Product Name         | Anti-KCNH1 Antibody Picoband™   |
|----------------------|---|
| Reactive Species     | Human, Mouse, Rat   |
| Description          | Boster Bio Anti-KCNH1 Antibody Picoband™ catalog # A01036-2. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.  |
| Application          | WB  |
| Clonality            | Polyclonal  |
| Formulation          | Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .  |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host                 | Rabbit  |
| Uniprot ID           | O95259  |

### **Technical Details**

| Immunogen                     | A synthetic peptide corresponding to a sequence at the C-terminus of human KCNH1, which shares 94.6% amino acid (aa) sequence identity with both mouse and rat KCNH1. |
|-------------------------------|---|
| Predicted Reactive Species    | Human   |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.   |
| Cross Reactivity              | No cross-reactivity with other proteins.  |
| Isotype                       | Rabbit IgG  |
|                               |   |





888-466-3604 | support@bosterbio.com | www.bosterbio.com

| Form                | Lyophilized  |
|---------------------|--|
| Concentration       | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.  |
| Purification        | Immunogen affinity purified.   |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Western blot, 0.1-0.5ug/ml |



## Anti-KCNH1 Antibody Picoband™ (A01036-2) Images




Figure 1. Western blot analysis of KCNH1 using anti-KCNH1 antibody (A01036-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human COLO-320 cell lysate,

Lane 2: human HepG2 cell lysate,

Lane 3: human A549 cell lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-KCNH1 antigen affinity purified polyclonal antibody (Catalog # A01036-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for KCNH1 at approximately 150KD. The expected band size for KCNH1 is at 111KD.

# Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-KCNH1 Antibody ™