

Anti-Tyrosine-protein kinase HCK Hck Antibody

Catalog Number: A01073-2

About HCK

Involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. May represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. Its activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity. Involved in osmotic stress-dependent SNCA 'Tyr-125' phosphorylation.

Gluck SL, et al. (2004) J Clin Invest; 114(12): 1696-1699

Benzing T, et al. (2001) Proc Natl Acad Sci U S A; 98(17): 9784-9789

Tian D, et al. (2002) Mol Cell Biol; 22(8): 2650-2662

Lu Z, et al. (2001) Mol Cell Biol; 21(12): 4016-4031

Krishnan HH, et al. (2006) J Virol; 80(3): 1167-1180

Overview

Product Name	Anti-Tyrosine-protein kinase HCK Hck Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Tyrosine-protein kinase HCK Hck Antibody catalog # A01073-2. Tested in ELISA, IF, IHC applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, IF, IHC
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P08631

Technical Details

Immunogen	Synthesized peptide derived from human Hck around the non-phosphorylation site of Y410.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG

Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:5000</p>

Anti-Tyrosine-protein kinase HCK Hck Antibody (A01073-2) Images




Figure 1. Western blotting validation for Anti-Tyrosine-protein kinase HCK Hck Antibody A01073-2

Western Blot (WB) analysis of specific cells using Hck polyclonal antibody. Electrophoresis was performed on a SDS-PAGE gel. To determine SDS-PAGE gel concentration

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Tyrosine-protein kinase HCK Hck Antibody