

Anti-Thymic stromal lymphopoietin TSLP Antibody

Catalog Number: A01096-2

About TSLP

Thymic stromal lymphopoietin (TSLP) has recently been identified as an important factor capable of driving dendritic cell maturation and activation. TSLP is a four-helix-bundle cytokine that is expressed mainly by barrier epithelial cells and is a potent activator of several cell types such as myeloid dendritic cells. TSLP is involved in the positive selection of regulatory T cells, maintenance of peripheral CD4+ T cell homeostasis and the induction of CD4+ T cell-mediated allergic reaction. TSLP is also capable of supporting the growth of fetal liver and adult B cell progenitors and their differentiation to the IgM-positive stage of B cell development. Amino acid sequence analysis has shown poor homology between human and mouse TSLP although they exhibit similar biological functions and are expressed in similar tissues. At least two differentially spliced isoforms of TSLP are known to exist.

Overview

Product Name	Anti-Thymic stromal lymphopoietin TSLP Antibody
Reactive Species	Human
Description	Boster Bio Anti-Thymic stromal lymphopoietin TSLP Antibody (Catalog # A01096-2). Tested in ELISA, WB, IHC-P applications. This antibody reacts with Human.
Application	ELISA, IHC-P, WB
Clonality	Polyclonal
Formulation	TSLP Antibody is supplied in PBS containing 0.02% sodium azide.
Storage Instructions	TSLP antibody can be stored at 4°C for three months and -20°C, stable for up to one year. Avoid repeated freeze-thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Host	Rabbit
Uniprot ID	Q969D9

Technical Details


Immunogen	Anti-TSLP antibody was raised against a peptide corresponding to 17 amino acids near the carboxy terminus of human TSLP. The immunogen is located within the last 50 amino acids of TSLP.
Predicted Reactive Species	Bovine, Pig
Isotype	IgG
Form	Liquid
Concentration	1 mg/mL
Purification	TSLP Antibody is affinity chromatography purified via peptide column.

Suggested Dilutions


WB: 0.25 - 2 ug/mL; IHC-P: 2.5 ug/mL.

Antibody validated: Western Blot in human samples; Immunohistochemistry and Immunocytochemistry in human samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher.


Anti-Thymic stromal lymphopoietin TSLP Antibody (A01096-2) Images




Western Blot Validation with Recombinant Protein Loading:
30 ng of human TSLP recombinant protein per lane.
Antibodies: TSLP A01096-2, 1h incubation at RT in 5%
NFDN/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate
at 1:10000 dilution. Lane 1: 0.25 ug/mL Lane 2: 0.5 ug/mL
Lane 3: 1 ug/mL



Western Blot Validation in Human Jurkat Cell Line Loading:
15 ug of lysates per lane. Antibodies: TSLP A01096-2 (A: 1
ug/mL, B: 2 ug/mL), 1h incubation at RT in 5% NFDN/TBST.
Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000
dilution.




Immunohistochemistry Validation of TSLP in Human Brain
Tissue Immunohistochemical analysis of paraffin-embedded
Human Brain Tissue using anti-TSLP antibody (A01096-2) at
2.5 ug/ml. Tissue was fixed with formaldehyde and blocked
with 10% serum for 1 h at RT; antigen retrieval was by heat
mediation with a citrate buffer (pH6). Samples were
incubated with primary antibody overnight at 4°C. A goat
anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary.
Counter stained with Hematoxylin.




Induced Expression Validation in Human Oral Keratinocytes
(Bjerkkan et al., 2015) TSLP expression detected by anti-TSLP
antibodies (A01096-2) was upregulated in cultured human
oral keratinocytes after 24 h in response to poly(I:C),


Regulated Expression Validation of TSLP in COPD Patients.
(Anzalone et al., 2018) (A) shows NHBE cells, in the absence
(Lane 2) or presence (Lane 3) of Tiotropium, were
stimulated with rhIL-17A. (B) shows NHBE cells, with ISs from
COPD patients untreated (Lane 2 and Lane 3) or treated
(Lane 4) with anti-IL-17A antibody. (A) and (B) show TSLP
expression decreases with the treatment of anti-cholinergic




drugs.



Induced Expression Validation of TSLP in Human Corneal Epithelium (Lin et al., 2018) Immunohistochemical images showing the TSLP protein detected by anti-TSLP antibodies in donor corneal tissues without (Control) or after exposure to IL-32 ex vivo; an isotype IgG antibody was used as a negative control. The production of TSLP was increased after IL-32 treatment.



Induced Expression Validation of TSLP in Human Corneal Tissues (Lin et al., 2013) Immunohistochemical images showing TSLP protein detected by anti-TSLP antibodies in ex vivo donor human corneal tissues without (B1) or with exposure to IL-33 (B2). An isotype IgG antibody (B3) was used as a negative control. Magnification 400X. The staining confirms the increased level of TSLP after IL-33 treatment.



Induced Expression Validation of TSLP in Mesenchymal Stromal Cells (MSCs). (Allakhverdi et al., 2013) Immunocytochemical images showing TSLP protein detected by anti-TSLP antibodies in BM-MSCs unstimulated (control) or stimulated with IL-1/TNF or supernatants of activated MNCs (MCactSNT). Isotype is used as a negative control. The production of TSLP increased with the stimulated conditions.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Thymic stromal lymphopoietin TSLP Antibody

For Research Use Only. Not for use in diagnostic procedures.