

Anti-CD40L/CD40LG Antibody Picoband™

Catalog Number: A01114-3

About CD40LG

CD40 ligand (CD40L) is a type II membrane protein of 261 amino acids on activated T cells that induces B cell proliferation and immunoglobulin secretion. It has homology with tumour necrosis factor- α and - β , and has important functions in B-cell activation and differentiation. Human CD40L with 5 exons, is mapped to the proximal region of the mouse X chromosome on Xq26.3-27.1, and can be detected on T cells but is absent from B cells and monocytes. Since CD40L is expressed on platelets and released from them on activation, its predictive value as a marker for clinical outcome and the therapeutic effect of inhibition of glycoprotein IIb/IIIa receptor in patients with acute coronary syndromes was investigated. The soluble CD40L may be involved in the process of restenosis and that it exerts its effect by triggering a complex group of inflammatory reactions on endothelial and mononuclear cells. CD40L plays a central role in the pathophysiology of acute coronary syndromes, and has a role in the pathogenesis of coronary artery lesions.

Overview

| | |
|----------------------|---|
| Product Name | Anti-CD40L/CD40LG Antibody Picoband™ |
| Reactive Species | Human, Mouse, Rat |
| Description | Boster Bio Anti-CD40L/CD40LG Antibody Picoband™ catalog # A01114-3. Tested in WB applications. This antibody reacts with Human, Mouse, Rat. |
| Application | WB |
| Clonality | Polyclonal |
| Formulation | Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | P29965 |

Technical Details

| | |
|-------------------------------|---|
| Immunogen | A synthetic peptide corresponding to a sequence in the middle region of human CD40L/CD40LG, which shares 82.8% and 79.3% amino acid (aa) sequence identity with mouse and rat CD40L/CD40LG, respectively. |
| Predicted Reactive Species | Human |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity | No cross-reactivity with other proteins. |

| | |
|---------------------|---|
| Isotype | Rabbit IgG |
| Form | Lyophilized |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. |
| Purification | Immunogen affinity purified. |
| Suggested Dilutions | <p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml</p> |

Anti-CD40L/CD40LG Antibody Picoband™ (A01114-3) Images




Figure 1. Western blot analysis of CD40LG using anti-CD40LG antibody (A01114-3).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human HL-60 whole cell lysates, Lane 2: human U-87MG whole cell lysates, Lane 3: rat liver tissue lysates, Lane 4: rat stomach tissue lysates, Lane 5: mouse spleen tissue lysates, Lane 6: mouse thymus tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD40LG antigen affinity purified polyclonal antibody (Catalog # A01114-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CD40LG at approximately 36KD. The expected band size for CD40LG is at 29KD.

1 Publications Citing This Product

1. PubMed ID: 19204010, Yang G, Fang Z, Liu Y, Zhang H, Shi X, Ji Q, Lin Q, Lin R. Evid Based Complement Alternat Med. 2011;2011:930324. Doi: 10.1093/Ecam/Nep013. Epub 2011 Jun 5. Protective Effects Of Chinese Traditional Medicine Buyang Huanwu Decoction On Myocardial In...

Visit bosterbio.com/anti-cd40l-cd40lg-picoband-trade-antibody-a01114-3-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD40L/CD40LG Antibody™