

Anti-TRAP/CD40L/Cd40lg Antibody Picoband® APC Conjugated

Catalog Number: A01114-5-APC

About Cd40lg

CD40 ligand(CD40L) is a type II membrane protein of 261 amino acids on activated T cells that induces B cell proliferation and immunoglobulin secretion. It has homology with tumour necrosis factor-alpha and -beta, and has important functions in B-cell activation and differentiation. Human CD40L with 5 exons, is mapped to the proximal region of the mouse X chromosome on Xq26.3-27.1, and can be detected on T cells but is absent from B cells and monocytes. Since CD40L is expressed on platelets and released from them on activation, its predictive value as a marker for clinical outcome and the therapeutic effect of inhibition of glycoprotein IIb /IIIa receptor in patients with acute coronary syndromes was investigated. The soluble CD40L may be involved in the process of restenosis and that it exerts its effect by triggering a complex group of inflammatory reactions on endothelial and mononuclear cells. CD40L plays a central role in the pathophysiology of acute coronary syndromes, and has a role in the pathogenesis of coronary artery lesions.

Overview

| | |
|----------------------|--|
| Product Name | Anti-TRAP/CD40L/Cd40lg Antibody Picoband® APC Conjugated |
| Reactive Species | Mouse, Rat |
| Application | Recommended applications are based on the parent unconjugated antibody (ELISA, IHC, WB). Customers may select suitable applications according to their experimental needs. |
| Clonality | Polyclonal |
| Formulation | Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ . |
| Storage Instructions | At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light. |
| Host | Rabbit |
| Uniprot ID | Q9Z2V2 |

Technical Details

| | |
|------------------|---|
| Immunogen | E.coli-derived rat TRAP/CD40L/Cd40lg recombinant protein (Position: M112-L260). |
| Cross Reactivity | No cross-reactivity with other proteins. |
| Isotype | Rabbit IgG |
| Form | Liquid |
| Concentration | 0.5 mg/mL |
| Purification | Immunogen affinity purified. |
| Conjugate | APC Excitation Wavelength: 633-647 nm |

| | |
|---------------------|--|
| | Emission Wavelength: 660 nm |
| Suggested Dilutions | Optimal dilutions should be determined by end users. |

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TRAP/CD40L/Cd40Ig Antibody - APC

For Research Use Only. Not for use in diagnostic procedures.