

## Anti-p75 NGF Receptor/NGFR Antibody Picoband®

Catalog Number: A01187

### About NGFR

The low-affinity nerve growth factor receptor (nerve growth factor receptor (TNFR superfamily, member 16), also called the LNGFR or p75 neurotrophin receptor) is one of the two receptor types for the neurotrophins, a family of protein growth factors that stimulate neuronal cells to survive and differentiate. LNGFR is a member of the tumor necrosis factor receptor (TNF receptor) superfamily indeed, LNGFR was the first member of this large family of receptors to be characterized. It is mapped to 17q21.33. Nerve growth factor receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain.

### Overview


Product Name	Anti-p75 NGF Receptor/NGFR Antibody Picoband®
Reactive Species	Human, Rat
Description	Boster Bio Anti-p75 NGF Receptor/NGFR Antibody Picoband® catalog # A01187. Tested in ELISA, Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IF, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P08138

### Technical Details


Immunogen	E.coli-derived human p75 NGF Receptor/NGFR recombinant protein (Position: K29-Q240).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Human, Rat Immunocytochemistry/Immunofluorescence, 2ug/ml, Human Flow Cytometry (Fixed), 1-3ug/1x10 <sup>6</sup> cells, Human ELISA, 0.1-0.5ug/ml, -


## Anti-p75 NGF Receptor/NGFR Antibody Picoband® (A01187) Images



Western blot analysis of p75 NGF Receptor/NGFR using anti-p75 NGF Receptor/NGFR antibody (A01187). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: human Sw620 whole cell lysates, Lane 3: human Caco-2 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-p75 NGF Receptor/NGFR antigen affinity purified polyclonal antibody (Catalog # A01187) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for p75 NGF Receptor/NGFR at approximately 45KD,75KD. The expected band size for p75 NGF Receptor/NGFR is at 45KD,75KD.



IF analysis of p75 NGF Receptor/NGFR using anti-p75 NGF Receptor/NGFR antibody (A01187). p75 NGF Receptor/NGFR was detected in immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 2ug/mL rabbit anti-p75 NGF Receptor/NGFR Antibody (A01187) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Flow Cytometry analysis of A431 cells using anti-p75 NGF Receptor/NGFR antibody (A01187). Overlay histogram showing A431 cells stained with A01187 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-p75 NGF Receptor/NGFR Antibody (A01187, 1ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

1. PubMed ID: 10.1007/s11302-015-9445-8, Up-regulation of P2X7 receptors mediating proliferation of Schwann cells after sciatic nerve injury

2. PubMed ID: 10.1021/acsbiomaterials.0c01474, Synthesis of Double Interfering Biodegradable Nano-MgO Micelle Composites and Their Effect on Parkinson's Disease

3. PubMed ID: 25666351, Chen Q, Zhang Z, Liu J, He Q, Zhou Y, Shao G, Sun X, Cao X, Gong A, Jiang P. Mol Cells. 2015 Mar;38(3):221-8. Doi: 10.14348/Molcells.2015.2170. Epub 2015 Feb 4. A Fibrin Matrix Promotes The Differentiation Of Emscs Isolated From Nasal Respiratory ...

Visit [bosterbio.com/anti-p75-ngf-receptor-ngfr-picoband-trade-antibody-a01187-boster.html](http://bosterbio.com/anti-p75-ngf-receptor-ngfr-picoband-trade-antibody-a01187-boster.html) to see all 3 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-p75 NGF Receptor/NGFR Antibody

For Research Use Only. Not for use in diagnostic procedures.