

## Anti-Choline Acetyltransferase/CHAT Antibody Picoband®

Catalog Number: A01192-3

### About CHAT

Choline acetyltransferase (commonly abbreviated as ChAT, but sometimes CAT) is a transferase enzyme responsible for the synthesis of the neurotransmitter acetylcholine. In humans, the choline acetyltransferase enzyme is encoded by the CHAT gene. This gene product is a characteristic feature of cholinergic neurons, and changes in these neurons may explain some of the symptoms of Alzheimer's disease. Polymorphisms in this gene have been associated with Alzheimer's disease and mild cognitive impairment. Mutations in this gene are associated with congenital myasthenic syndrome associated with episodic apnea. Multiple transcript variants encoding different isoforms have been found for this gene, and some of these variants have been shown to encode more than one isoform.

### Overview


Product Name	Anti-Choline Acetyltransferase/CHAT Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Choline Acetyltransferase/CHAT Antibody Picoband® catalog # A01192-3. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P28329

### Technical Details


Immunogen	E.coli-derived human CHAT recombinant protein (Position: D446-R652). Human CHAT shares 88.4% and 87% amino acid (aa) sequence identity with mouse and rat CHAT, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Mouse, Rat, Human Western blot, 0.1-0.5ug/ml, Mouse, Rat, Human


## Anti-Choline Acetyltransferase/CHAT Antibody Picoband® (A01192-3) Images



IHC analysis of CHAT using anti-CHAT antibody (A01192-3). CHAT was detected in paraffin-embedded section of mouse brain tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CHAT Antibody (A01192-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



IHC analysis of CHAT using anti-CHAT antibody (A01192-3). CHAT was detected in paraffin-embedded section of rat brain tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CHAT Antibody (A01192-3) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



Western blot analysis of CHAT using anti-CHAT antibody (A01192-3). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat testis tissue lysates, Lane 2: mouse testis tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CHAT antigen affinity purified polyclonal antibody (Catalog # A01192-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CHAT at approximately 83KD. The expected band size for CHAT is at 83KD.

## 8 Publications Citing This Product

1. PubMed ID: 10.1111/j.1440-1681.2009.05149.x, PROTECTIVE EFFECTS OF ICARIIN ON COGNITIVE DEFICITS INDUCED BY CHRONIC CEREBRAL HYPOPERFUSION IN RATS
2. PubMed ID: 26509167, Overexpression of NTRK1 Promotes Differentiation of Neural Stem Cells into Cholinergic Neurons

3. PubMed ID: 18983683, Chemokine CXCL12 and its receptor CXCR4 expression are associated with perineural invasion of prostate cancer

Visit [bosterbio.com/anti-chat-picoband-trade-antibody-a01192-3-boster.html](https://bosterbio.com/anti-chat-picoband-trade-antibody-a01192-3-boster.html) to see all 8 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Choline Acetyltransferase/CHAT Antibody

For Research Use Only. Not for use in diagnostic procedures.