

## Anti-MAP2 Antibody Picoband®

Catalog Number: A01201

### About MAP2

Microtubule-associated protein 2 is a protein that in humans is encoded by the MAP2 gene. This gene encodes a protein that belongs to the microtubule-associated protein family. The proteins of this family are thought to be involved in microtubule assembly, which is an essential step in neurogenesis. The products of similar genes in rat and mouse are neuron-specific cytoskeletal proteins that are enriched in dendrites, implicating a role in determining and stabilizing dendritic shape during neuron development. A number of alternatively spliced variants encoding distinct isoforms have been described.

### Overview

Product Name	Anti-MAP2 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-MAP2 Antibody Picoband® catalog # A01201. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P11137


### Technical Details

Immunogen	E. coli-derived human MAP2 recombinant protein (Position: Q64-L133).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Suggested Dilutions

Western blot, 0.1-0.5ug/ml  
ELISA, 0.1-0.5ug/ml

## Anti-MAP2 Antibody Picoband® (A01201) Images



Western blot analysis of MAP2 using anti-MAP2 antibody (A01201). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MAP2 antigen affinity purified polyclonal antibody (Catalog # A01201) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MAP2 at approximately 280KD. The expected band size for MAP2 is at 199KD.

## 20 Publications Citing This Product

1. PubMed ID: , The effect of acetylcholine-like biomimetic polymers on neuronal growth
2. PubMed ID: 33582188, Wu K, Yue J, Shen K, He J, Zhu G, Liu S, Yang H, Zhang C. Expression and cellular distribution of FGF13 in cortical tubers of the tuberous sclerosis complex. *Neurosci Lett*. 2021 Feb 11;135714. doi:10.1016/j.neulet.2021.135714. Epub ahead of print. PMID:33582188.
3. PubMed ID: 27853172, Lithium promotes DNA stability and survival of ischemic retinal neurocytes by upregulating DNA ligase IV

Visit [bosterbio.com/anti-map2-picoband-trade-antibody-a01201-boster.html](https://www.bosterbio.com/anti-map2-picoband-trade-antibody-a01201-boster.html) to see all 20 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MAP2 Antibody

For Research Use Only. Not for use in diagnostic procedures.