

Anti-TXN Antibody (C-term)

Catalog Number: A01219

About TXN

Participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. Plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity. Induces the FOS/JUN AP-1 DNA-binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity. ADF augments the expression of the interleukin-2 receptor TAC (IL2R/P55).

Overview

Product Name	Anti-TXN Antibody (C-term)
Reactive Species	Human
Description	Boster Bio Anti-TXN Antibody (C-term) (Catalog # A01219). Tested in IHC-P, WB application(s). This antibody reacts with Human.
Application	IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P10599


Technical Details

Immunogen	This TXN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 66-94 amino acids from the C-terminal region of human TXN.
Predicted Reactive Species	Bovine, Mouse, Pig, Rabbit, Rat
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.


Suggested Dilutions

WB: 1:1000
IHC-P: 1:25


Anti-TXN Antibody (C-term) (A01219) Images



TXN Antibody (C-term) western blot analysis in A549,MDA-MB453 cell line lysates (35ug/lane). This demonstrates the TXN antibody detected the TXN protein (arrow).



Immunohistochemical analysis of paraffin-embedded H. esophagus section using TXN Antibody (C-term) (Cat#A01219). A01219 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. stomach section using TXN Antibody (C-term) (Cat#A01219). A01219 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TXN Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.