

Anti-NUP98 Antibody Picoband®

Catalog Number: A01301-1

About NUP98

Nuclear pore complex protein Nup98-Nup96 is a protein that in humans is encoded by the NUP98 gene. This gene is one of several genes located in the imprinted gene domain of 11p15.5, an important tumor-suppressor gene region. NUP98 is a peripheral nucleoporin located at both the cytoplasmic and nuclear sides of the central channel of the NPC. NUP98 phosphorylation is critical for NPC disassembly at the onset of mitosis. It also plays roles in gene expression, mitotic checkpoint, and pathogenesis. Ligand blot analysis suggested that NUP98 can function as a docking protein for cytosol-mediated docking of import substrates. In addition to that, NUP98 is a target of the vesicular stomatitis virus M protein-mediated inhibition of mRNA nuclear export.

Overview


| | |
|----------------------|--|
| Product Name | Anti-NUP98 Antibody Picoband® |
| Reactive Species | Human, Monkey, Mouse, Rat |
| Description | Boster Bio Anti-NUP98 Antibody Picoband® catalog # A01301-1. Tested in WB, ICC/IF, FCM, ELISA applications. This antibody reacts with Human, Monkey, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |
| Application | ELISA, Flow Cytometry, IF, ICC, WB |
| Clonality | Polyclonal |
| Formulation | Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ . |
| Storage Instructions | At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing. |
| Host | Rabbit |
| Uniprot ID | P52948 |

Technical Details


| | |
|-------------------------------|---|
| Immunogen | E.coli-derived human NUP98 recombinant protein (Position: H728-I1787). Human NUP98 shares 89.1% and 89% amino acid (aa) sequence identity with mouse and rat NUP98, respectively. |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC. |
| Cross Reactivity | No cross reactivity with other proteins. |
| Isotype | IgG |

| | |
|---------------------|---|
| Form | Lyophilized |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml. |
| Purification | Immunogen affinity purified. |
| Suggested Dilutions | Western blot, 0.25-0.5 µg/ml, Human, Monkey, Mouse, Rat Immunocytochemistry/Immunofluorescence, 5 µg/ml, Human Flow Cytometry (Fixed), 1-3 µg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 µg/ml, - |


Anti-NUP98 Antibody Picoband® (A01301-1) Images



Western blot analysis of NUP98 using anti-NUP98 antibody (A01301-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Caco-2 whole cell lysates, Lane 2: human HeLa whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: monkey COS-7 whole cell lysates, Lane 5: rat NRK whole cell lysates, Lane 6: rat PC-12 whole cell lysates, Lane 7: mouse RAW264.7 whole cell lysates, Lane 8: mouse NIH/3T3 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NUP98 antigen affinity purified polyclonal antibody (Catalog # A01301-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for NUP98 at approximately 98,105 kDa. The expected band size for NUP98 is at 198 kDa.



IF analysis of NUP98 using anti-NUP98 antibody (A01301-1). NUP98 was detected in an immunocytochemical section of A549 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-NUP98 Antibody (A01301-1) overnight at 4°C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Flow Cytometry analysis of K562 cells using anti-NUP98 antibody (A01301-1). Overlay histogram showing K562 cells stained with A01301-1 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-NUP98 Antibody (A01301-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NUP98 Antibody

For Research Use Only. Not for use in diagnostic procedures.