

Anti-CB2 Antibody (C-term)

Catalog Number: A01307-3

About CNR2

The cannabinoid delta-9-tetrahydrocannabinol is the principal psychoactive ingredient of marijuana. The proteins encoded by this gene and the cannabinoid receptor 1 (brain) (CNR1) gene have the characteristics of a guanine nucleotide-binding protein (G-protein)-coupled receptor for cannabinoids. They inhibit adenylate cyclase activity in a dose-dependent, stereoselective, and pertussis toxin-sensitive manner. These proteins have been found to be involved in the cannabinoid-induced CNS effects (including alterations in mood and cognition) experienced by users of marijuana. The cannabinoid receptors are members of family 1 of the G-protein-coupled receptors.

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview


Product Name	Anti-CB2 Antibody (C-term)
Reactive Species	Human
Description	Boster Bio Anti-CB2 Antibody (C-term) (Catalog # A01307-3). Tested in WB, IHC-P, Flow Cytometry application(s). This antibody reacts with Human.
Application	Flow Cytometry, IHC-P, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P34972

Technical Details


Immunogen	This CB2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 329-356 amino acids from the C-terminal region of human CB2.
Predicted Reactive Species	Bovine, Xenopus
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western

	blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:1000 IHC-P: 1:10-1:50 FC: 1:25


Anti-CB2 Antibody (C-term) (A01307-3) Images




CB2 Antibody (C-term) western blot analysis in A431 cell line lysates (35ug/lane). This demonstrates the CB2 antibody detected the CB2 protein (arrow).



Immunohistochemical analysis of paraffin-embedded R. brain section using CB2 Antibody (C-term). A01307-3 was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.





Immunohistochemical analysis of paraffin-embedded H. brain section using CB2 Antibody (C-term). A01307-3 was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



CB2 antibody (C-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human skin carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CB2 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

CB2 Antibody (C-term) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Flow cytometric analysis of Jurkat cells using CB2 Antibody (C-term) compared to an isotype control of rabbit IgG (blue). A01307-3 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

Anti-CB2 Antibody (C-term)

For Research Use Only. Not for use in diagnostic procedures.