

Anti-Human IDE DyLight® 488 conjugated Antibody

Catalog Number: A01358-Dyl488

About IDE

Insulin-degrading enzyme, also known as IDE, is an enzyme. This gene encodes a zinc metallopeptidase that degrades intracellular insulin, and thereby terminates insulins activity, as well as participating in intercellular peptide signalling by degrading diverse peptides such as glucagon, amylin, bradykinin, and kallidin. The preferential affinity of this enzyme for insulin results in insulin-mediated inhibition of the degradation of other peptides such as beta-amyloid. Deficiencies in this protein's function are associated with Alzheimer's disease and type 2 diabetes mellitus but mutations in this gene have not been shown to be causitive for these diseases. This protein localizes primarily to the cytoplasm but in some cell types localizes to the extracellular space, cell membrane, peroxisome, and mitochondrion. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Overview

Product Name	Anti-Human IDE DyLight® 488 conjugated Antibody
Reactive Species	Human
Description	Boster Bio Anti-Human IDE DyLight® 488 conjugated Antibody catalog # A01358-Dyl488. Tested in Flow Cytometry applications. This antibody reacts with Human.
Conjugate	DyLight®488
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na2HPO4, 0.02% NaN3.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P14735

Technical Details

Immunogen	E. coli-derived human IDE recombinant protein (Position: F485-K756).
Predicted Reactive Species	Human
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

	procedure.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Flow Cytometry, 1-3ug/1x10 ⁶ cells



Anti-Human IDE DyLight® 488 conjugated Antibody (A01358-Dyl488) Images



Boster Kit Box

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Human IDE DyLight® 488 conjugated Antibody