

Anti-Fos B/FOSB Antibody

Catalog Number: A01569-1

About FOSB

FOSB, FBJ murine osteosarcoma viral oncogene homolog B, is a protein that, in humans, is encoded by the FOSB gene. FOSB is a member of Fos gene family which consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. The FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. The FOSB gene is mapped to 19q13.32. Delta FOSB is a truncated splice variant of FOSB. Delta FosB has been implicated in the development of drug addiction and control of the reward system in the brain, and is linked to changes in a number of other gene products such as CREB and sirtuins. Delta FosB also regulates the commitment of mesenchymal precursor cells to the adipocyte or osteoblast lineage.

Overview

Product Name	Anti-Fos B/FOSB Antibody
Reactive Species	Human
Description	Boster Bio Anti-Fos B/FOSB Antibody catalog # A01569-1. Tested in ELISA, IF, IHC, ICC applications. This antibody reacts with Human.
Application	ELISA, IF, IHC, ICC
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P53539

Technical Details

Immunogen	E.coli-derived human FOSB recombinant protein (Position: M1-S332).
Recommended Detection Systems	Boster recommends HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human Immunocytochemistry/Immunofluorescence, 5ug/ml, Human ELISA, 0.1-0.5ug/ml, -

Anti-Fos B/FOSB Antibody (A01569-1) Images




Figure 1. IHC analysis of Fos B/FOSB using anti-Fos B/FOSB antibody (A01569-1). Fos B/FOSB was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Fos B/FOSB Antibody (A01569-1) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




Figure 2. IHC analysis of Fos B/FOSB using anti-Fos B/FOSB antibody (A01569-1). Fos B/FOSB was detected in a paraffin-embedded section of human gallbladder adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Fos B/FOSB Antibody (A01569-1) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




Figure 3. IHC analysis of Fos B/FOSB using anti-Fos B/FOSB antibody (A01569-1). Fos B/FOSB was detected in a paraffin-embedded section of human gallbladder adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Fos B/FOSB Antibody (A01569-1) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




Figure 4. IHC analysis of Fos B/FOSB using anti-Fos B/FOSB antibody (A01569-1). Fos B/FOSB was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Fos B/FOSB Antibody (A01569-1) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex

(SABC) (Catalog # SA1022) with DAB as the chromogen.




Figure 5. IF analysis of Fos B/FOSB using anti-Fos B/FOSB antibody (A01569-1).

Fos B/FOSB was detected in an immunocytochemical section of SiHa cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-Fos B/FOSB Antibody (A01569-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Fos B/FOSB Antibody