

Anti-CYP7A1 Antibody Picoband®

Catalog Number: A01601

About CYP7A1

CYP7A1 (Cytochrome P450 Subfamily VIIA Polypeptide 1), also called CYP7 or CHOLESTEROL 7-ALPHA-HYDROXYLASE, is an enzyme that in humans is encoded by the CYP7A1 gene. Using both mouse-human somatic cell hybrids and in situ chromosomal hybridization, Cohen et al. (1992) mapped the CYP7 gene to 8q11-q12. By transfection of reporter constructs, mutation analysis, and DNase footprinting, Molowa et al. (1992) identified areas of the CYP7A1 promoter region that showed hepatocyte-specific activation. They found HNF3 to be an activator of CYP7A1 activity. Agellon et al. (2002) found that wildtype mice and mice transgenic for human CYP7A1 respond differently to cholesterol feeding. Cholesterol feeding stimulated Cyp7a1 mRNA abundance and enzymatic activity in wildtype mice, but repressed human CYP7A1 mRNA and activity in transgenic mice.

Overview


Product Name	Anti-CYP7A1 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-CYP7A1 Antibody Picoband® catalog # A01601. Tested in ELISA, Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, Flow Cytometry, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P22680

Technical Details


Immunogen	E.coli-derived human CYP7A1 recombinant protein (Position: I10-L504).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5ug/ml, Human Immunohistochemistry(Paraffin-embedded Section, 2-5 ug/ml, Human Immunocytochemistry/Immunofluorescence, 5 ug/ml, Human Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Human ELISA, 0.1-0.5ug/ml, -


Anti-CYP7A1 Antibody Picoband® (A01601) Images



Western blot analysis of CYP7A1 using anti-CYP7A1 antibody (A01601). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HCCT tissue lysates, Lane 2: human HepG2 whole cell lysates, Lane 3: human HL-60 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CYP7A1 antigen affinity purified polyclonal antibody (Catalog # A01601) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CYP7A1 at approximately 58 kDa. The expected band size for CYP7A1 is at 58 kDa.




IHC analysis of CYP7A1 using anti-CYP7A1 antibody (A01601). CYP7A1 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CYP7A1 Antibody (A01601) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Flow Cytometry analysis of SiHa cells using anti-CYP7A1 antibody (A01601). Overlay histogram showing SiHa cells stained with A01601 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CYP7A1 Antibody (A01601, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

IF analysis of CYP7A1 using anti-CYP7A1 antibody (A01601). CYP7A1 was detected in an immunocytochemical section of SiHa cells. Enzyme antigen retrieval was performed using



IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 4 ug/mL rabbit anti-CYP7A1 Antibody (A01601) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

1 Publications Citing This Product

1. PubMed ID: 10.1093/toxsci/kfv155, Cellular Accumulation and Toxic Effects of Bile Acids in Cyclosporine A-Treated HepaRG Hepatocytes

Visit bosterbio.com/anti-cyp7a1-picoband-trade-antibody-a01601-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CYP7A1 Antibody

For Research Use Only. Not for use in diagnostic procedures.