

Anti-Human CTNNA1 DyLight® 488 conjugated Antibody

Catalog Number: A01617-Dyl488

About CTNNA1

CTNNA1, also known as Catenin alpha-1 or Catenin (cadherin-associated protein), alpha 1, is a protein that in humans is encoded by the CTNNA1 gene. It is mapped to 5q31.2. When surface epithelium CTNNA1 was ablated, hair follicle development was blocked and epidermal morphogenesis was dramatically affected, with defects in adherens junction formation, intercellular adhesion, and epithelial polarity. In vitro, CTNNA1 null keratinocytes were poorly contact inhibited and grew rapidly. These differences were not dependent upon intercellular adhesion and were in marked contrast to keratinocytes conditionally null for another essential intercellular adhesion protein, desmoplakin. Knockout keratinocytes exhibited sustained activation of the Ras-MAPK cascade due to aberrations in growth factor responses. It is concluded that features of precancerous lesions often attributed to defects in cell cycle regulatory genes can be generated by compromising the function of CTNNA1.

Overview

Product Name	Anti-Human CTNNA1 DyLight® 488 conjugated Antibody
Reactive Species	Human
Description	Boster Bio Anti-Human CTNNA1 DyLight® 488 conjugated Antibody catalog # A01617-Dyl488. Tested in Flow Cytometry applications. This antibody reacts with Human.
Conjugate	DyLight®488
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P35221

Technical Details

Immunogen	E.coli-derived human CTNNA1 recombinant protein (Position: D143-D292). Human CTNNA1 shares 98% amino acid (aa) sequence identity with mouse CTNNA1.
Predicted Reactive Species	Human
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid

Concentration	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Flow Cytometry, 1-3ug/1x10⁶ cells</p>

Anti-Human CTNNA1 DyLight® 488 conjugated Antibody (A01617-Dyl488) Images




Figure 1. Flow Cytometry analysis of Hela cells using anti-Human CTNNA1 antibody (A01617-Dyl488). Overlay histogram showing Hela cells stained with A01617-Dyl488 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Human CTNNA1 Antibody (A01617-Dyl488, 1 μ g/1 \times 10⁶ cells) for 30 min at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 μ g/1 \times 10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

2 Publications Citing This Product

1. PubMed ID: 15484291, Prognostic and clinicopathological features of E-cadherin, β -catenin, γ -catenin and cyclin D1 expression in human esophageal squamous cell carcinoma
2. PubMed ID: 27065079, TALENs-directed knockout of the full-length transcription factor Nrf1 that represses malignant behaviour of human hepatocellular carcinoma (HepG2) cells

Visit bosterbio.com/anti-human-ctnna1-dylight-488-conjugated-antibody-a01617-dyl488-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Human CTNNA1 DyLight® 488 conjugated Antibody