

Anti-PGD Antibody Picoband® APC Conjugated

Catalog Number: A01623-3-APC

About PGD

6-phosphogluconate dehydrogenase is the second dehydrogenase in the pentose phosphate shunt. Deficiency of this enzyme is generally asymptomatic, and the inheritance of this disorder is autosomal dominant. Hemolysis results from combined deficiency of 6-phosphogluconate dehydrogenase and 6-phosphogluconolactonase suggesting a synergism of the two enzymopathies. Several transcript variants encoding different isoforms have been found for this gene.

Overview

Product Name	Anti-PGD Antibody Picoband® APC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, IF, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P52209

Technical Details

Immunogen	E.coli-derived human PGD recombinant protein (Position: D24-R447). Human PGD shares 93.9% and 92.9% amino acid (aa) sequence identity with mouse and rat PGD, respectively.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	APC Excitation Wavelength: 633-647 nm Emission Wavelength: 660 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PGD Antibody - APC

For Research Use Only. Not for use in diagnostic procedures.