

Anti-CaM Kinase IV CAMK4 Antibody

Catalog Number: A01905

About CAMK4

CaM Kinase IV (also known as CAM kinase-GR and CaMK IV) is a calcium/ calmodulin-dependent protein kinase belonging to a proposed calcium-triggered signaling cascade. This kinase may be involved in the transcriptional regulation of microtubule dynamics. In vitro, CaMK IV phosphorylates CREB1, CREBBP, PRM2, MEF2A, MEF2D and STMN1/OP18. CaMK IV may also be involved in spermatogenesis and may play a role in the consolidation/ retention of hippocampus-dependent long-term memory. CaMK IV must be phosphorylated to be maximally active and is phosphorylated by CAMKK1 or CAMKK2. In addition autophosphorylation of the N-terminus is required for full activation. Autophosphorylation of Ser-336 allows the kinase to switch to a Ca(2+)/calmodulin-independent state. Most likely the kinase is inactivated by the serine/ threonine protein phosphatase 2A. CaMK IV is a monomer that is located within the cytoplasm and nucleus and substantial localization occurs in certain neuronal nuclei. In spermatids CaMK IV is associated with chromatin and the nuclear matrix. CaMK IV is also specifically expressed in epithelial ovarian cancer tissue.

Overview


Product Name	Anti-CaM Kinase IV CAMK4 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-CaM Kinase IV CAMK4 Antibody (Catalog # A01905). Tested in ELISA, IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at 4°C prior to restoration. For extended storage aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship at ambient temperature.)
Host	Rabbit
Uniprot ID	Q16566

Technical Details

Immunogen	This antiserum was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near amino acids 300-325 of Human CaM Kinase IV protein.
Predicted Reactive Species	Bovine, Canine, Chimpanzee
Cross Reactivity	No cross reactivity with other proteins.

Isotype	Antiserum
Form	Lyophilized
Concentration	70 mg/mL by Refractometry
Purification	This antiserum is directed against human CaM Kinase IV protein. The product was delipidated, defibrinated followed by buffering and clarification. A BLAST analysis was used to suggest reactivity with this protein from human, mouse, and rat based on 100% homology for the immunogen sequence. Cross-reactivity with CaM Kinase IV homologues from other sources has not been determined.
Suggested Dilutions	ELISA: 1:5,000 - 1:25,000 IHC: 1:500 WB: 1:500 - 1:2,000 This antiserum has been tested for use in ELISA, IHC, and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band ~ 52 kDa in size corresponding to CaM Kinase IV by western blotting in the appropriate cell lysate or extract.

Anti-CaM Kinase IV CAMK4 Antibody (A01905) Images



Immunohistochemistry of Anti-CAMK4 antibody. Tissue: human brain cortex was formalin fixed and paraffin embedded. No pre-treatment of sample was required. Primary Antibody: Anti-CaM Kinase IV was diluted 1:500. The image shows the localization of antibody as the precipitated red signal, with a hematoxylin purple nuclear counter stain.



Western blot using Boster's Anti-CaM Kinase IV antibody. Lane 1: Rat Brain, adult, WCL . Lane 2: Jurkat Whole Cell Lysate . Lane 3: Rat Brain, adult, WCL , preincubated with immunizing peptide. Lane 4: Jurkat Whole Cell Lysate , preincubated with immunizing peptide. Load: 35µg lysate/lane. Primary Antibody: Anti-CaM Kinase IV at 1:1,000 for 2hr at RT. Secondary Antibody: Goat Anti-Rabbit IgG IRDye800 at 1:10,000 for 45 mins at RT. Block: Fluorescent Buffer 30 mins at RT. Results: band ~52 kDa corresponding to CaM Kinase IV (arrowhead). CaM Kinase IV was similarly detected on lysates from mouse brain (not shown). IRDye800 fluorescence image was captured using the Odyssey® Infrared Imaging System developed by LI-COR. IRDye is a trademark of LI-COR, Inc.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CaM Kinase IV CAMK4 Antibody

For Research Use Only. Not for use in diagnostic procedures.