

Anti-HGD Antibody Picoband® Cy3 Conjugated

Catalog Number: A01909-1-Cy3

About HGD

The HGD gene encodes homogentisate 1,2-dioxygenase (HGD), an enzyme involved in the catabolism of phenylalanine and tyrosine. This enzyme is involved in the catabolism of the amino acids tyrosine and phenylalanine. Mutations in this gene are the cause of the autosomal recessive metabolism disorder alkaptonuria. This gene is mapped to chromosome 3q21-q23 by a preliminary PCR screen of hamster/human somatic cell hybrid genomic DNA samples and by fluorescence in situ hybridization.

Overview

| | |
|----------------------|---|
| Product Name | Anti-HGD Antibody Picoband® Cy3 Conjugated |
| Reactive Species | Human, Mouse, Rat |
| Application | Recommended applications are based on the parent unconjugated antibody (ELISA, Flow Cytometry, IF, IHC, ICC, WB). Customers may select suitable applications according to their experimental needs. |
| Clonality | Polyclonal |
| Formulation | Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N. |
| Storage Instructions | At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light. |
| Host | Rabbit |
| Uniprot ID | Q93099 |

Technical Details

| | |
|---------------------|--|
| Immunogen | E. coli-derived human HGD recombinant protein (Position: D374-N445). |
| Cross Reactivity | No cross-reactivity with other proteins. |
| Isotype | Rabbit IgG |
| Form | Liquid |
| Concentration | 0.5 mg/mL |
| Purification | Immunogen affinity purified. |
| Conjugate | Cy3 Excitation Wavelength: 554 nm Emission Wavelength: 568 nm |
| Suggested Dilutions | Optimal dilutions should be determined by end users. |

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HGD Antibody - Cy3

For Research Use Only. Not for use in diagnostic procedures.