

Anti-HDAC-7 (N-terminus) Antibody

Catalog Number: A01913

About HDAC7

HDAC7 is a member of the class II mammalian histone deacetylases, which plays an important role in modulating the eukaryotic chromatin structure. Human HDAC7 is composed of 912 amino acid residues. Although HDAC7 is localized mostly to the cell nucleus, it is also found in the cytoplasm, suggesting nucleo-cytoplasmic shuttling. The histone deacetylase activity of HDAC7 maps to a carboxy-terminal domain and is dependent on interaction with class I HDACs in the nucleus. It is an active component of different transcriptional corepressor complexes that can be recruited to specific promoter regions via interactions with a growing number of sequence specific transcriptional factors. HDAC7 catalyzes removal of acetyl-groups from acetyl-lysines of histones and promotes compaction of chromatin in these regions, leading to the inhibition of gene transcription. Anti-HDAC7 antibodies are ideal for researches interested in Breast Cancer, Cancer, Cell Cycle and Replication, Chromatin Research, Epigenetics, and Histone Deacetylases research.

Overview


Product Name	Anti-HDAC-7 (N-terminus) Antibody
Reactive Species	Human, Mouse, Rat, Primate
Description	Boster Bio Anti-HDAC-7 (N-terminus) Antibody (Catalog # A01913). Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat, Primate.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is six (6) months from date of opening.
Host	Rabbit
Uniprot ID	Q8WUI4

Technical Details

Immunogen	HDAC7 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to the N terminal region of human HDAC7.
Predicted Reactive Species	Primate
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG

Form	Liquid (sterile filtered)
Concentration	1.16 mg/ml by UV absorbance at 280 nm
Purification	Anti-HDAC7 was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody is specific towards HDAC7. A BLAST analysis was used to suggest cross-reactivity with Human, Mouse, Rat and Primate based on 100% sequence homology. Cross-reactivity with HDAC7 from other sources has not been determined.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>ELISA: 1:20,000 - 1:60,000</p> <p>IHC: 1:100-1:500</p> <p>IF Microscopy: 1:100-1:500</p> <p>WB: 1 µg/ml</p>

Anti-HDAC-7 (N-terminus) Antibody (A01913) Images



Western blot analysis of HDAC-7 expression in mouse brain homogenate (lane 1) and mouse brain homogenate blocked with peptide (lane 2). HDAC-7 at 103KD was detected using rabbit anti-HDAC-7 Antigen Affinity purified polyclonal antibody (Catalog # A01913) at 1 ug/mL. Other band(s): HDAC7 isoforms ~70 and 40kDa. The blot was developed using chemiluminescence (ECL) method (Catalog # EK1002).

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HDAC-7 (N-terminus) Antibody