

Anti-Cul5/Cullin 5 Antibody

Catalog Number: A01925

About CUL5

Cullins assemble a potentially large number of ubiquitin ligases by binding to the RING protein ROC1 to catalyse polyubiquitination, as well as binding to various specificity factors to recruit substrates. Cullin 5 is a component of E3 ubiquitin ligase complexes, which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. Cullin 5 seems to be involved proteasomal degradation of p53/TP53 stimulated by adenovirus E1B-55 kDa protein. Cullin 5 may form a cell surface vasopressin receptor. Cullin 5 is part of a E3 ubiquitin ligase complex with elongin BC complex (TCEB1 and TCEB2), RBX1 and MUF1, complexes with elongin BC complex (TCEB1 and TCEB2), RXB1 and TCEB3 or SOCS1 or WSB1; elongin BC complex (TCEB1 and TCEB2), RBX1 and VHL; elongin BC complex (TCEB1 and TCEB2), RBX1, adenovirus type 5 E1B-55kDa protein and adenovirus type 5 E4-orf6. Cullin 5 Interacts with RBX1 and RNF7.

Overview


Product Name	Anti-Cul5/Cullin 5 Antibody
Reactive Species	Human
Description	Boster Bio Anti-Cul5/Cullin 5 Antibody (Catalog # A01925). Tested in IHC, IP, WB applications. This antibody reacts with Human.
Application	ELISA, IP, IHC, WB
Clonality	Polyclonal
Formulation	0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship on dry ice.)
Host	Rabbit
Uniprot ID	Q93034

Technical Details

Immunogen	This antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an N-Terminal region near amino acids 1-25 of Human Cul5 (N-terminus) coupled to KLH.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	Antiserum
Form	Liquid (sterile filtered)

Concentration	85 mg/mL by Refractometry
Purification	This product is monospecific antiserum processed by delipidation and defibrination followed by sterile filtration. This product reacts with human Cullin 5. Cross-reactivity is expected against mouse Cul5 based on a high degree of sequence homology. Cross-reactivity with other human cullins may occur.
Suggested Dilutions	ELISA: 1:2,000 - 1:10,000 IHC: User optimized IP: User optimized WB: 1:500 - 1:1,000 Anti-Cul5 has been tested by immunohistochemistry. This antibody reacts with human Cul5 by western blot and immunoprecipitation. The antibody immunoprecipitates protein from cell lysates (using 293T, U2OS and others). This antibody also co-immunoprecipitates associated proteins. A 90.9 kDa band corresponding to human Cul5 is detected. Most cell lines expressing Cul4A can be used as a positive control. Researchers should determine optimal titers for other applications.

Anti-Cul5/Cullin 5 Antibody (A01925) Images



Most modifiers mature by proteolytic processing from inactive precursors (a; amino acid). Arrowheads point to the cleavage sites. Ubiquitin is expressed either as polyubiquitin or as a fusion with ribosomal proteins. Conjugation requires activating (E1) and conjugating (E2) enzymes that form thioesters (S) with the modifiers. Modification of cullins by RUB involves SCF(SKP1/cullin-1/F-box protein) /CBC(cullin-2/elongin B/elonginC) -like E3 enzymes that are also involved in ubiquitination. In contrast to ubiquitin, the UBLs do not seem to form multi-UBL chains. UCRP(ISG15) resembles two ubiquitin moieties linked head-to-tail. Whether HUB1 functions as a modifier is currently unclear. APG12 and URM1 are distinct from the other modifiers because they are unrelated in sequence to ubiquitin. Data contributed by S.Jentsch, see references below.



Boster's Anti-CUL5 antibody was diluted 1:500 to detect CUL5 in human kidney tissue. Tissue was formalin fixed and paraffin embedded. No pre-treatment of sample was required. The image shows the localization of antibody as the precipitated red signal, with a hematoxylin purple nuclear counter stain.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cul5/Cullin 5 Antibody

For Research Use Only. Not for use in diagnostic procedures.