BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-OMI HTRA2 Antibody

Catalog Number: A01941

About HTRA2

Inhibitor of apoptosis proteins (IAPs) were initially identified in baculoviruses as proteins that inhibit apoptosis of the host cells to allow time for viral replication. Cellular homologues containing at least one baculoviral IAP repeat (BIR) motif essential for their anti-apoptosis activity have been identified in yeasts and higher organisms and often act by binding and inhibiting processed caspases. The activity of these proteins can be modulated by the expression of proteins such as Smac/DIABLO and XAF-1 which displace or prevent the binding of caspases by IAPs. Recently, a mitochondrial serine protease termed Omi/HtrA2 has been found to bind IAPs. Similar to Smac, Omi possesses a conserved IAP-binding motif, but acts to cleave IAPs to irreversibly inactivate IAPs and promote apoptosis.

Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-OMI HTRA2 Antibody
Reactive Species	Human
Description	Boster Bio Anti-OMI HTRA2 Antibody (Catalog # A01941). Tested in ELISA, WB, IHC-P, IF applications. This antibody reacts with Human.
Application	ELISA, IF, IHC-P, WB
Clonality	Polyclonal
Formulation	OMI Antibody is supplied in PBS containing ≤0.09% sodium azide.
Storage Instructions	OMI antibody can be stored at 4°C, stable for up to one year. Avoid repeated freeze-thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Host	Rabbit
Uniprot ID	O43464

Technical Details


Immunogen	OMI antibody was raised against a peptide corresponding to 16 amino acids near the C-terminus of human Omi. The immunogen is located within amino acids 70 - 120 of OMI.
Predicted Reactive Species	Bovine

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Isotype	IgG
Form	Liquid
Concentration	1 mg/mL
Purification	OMI Antibody is Ion exchange chromatography purified.
Suggested Dilutions	OMI antibody can be used for detection of OMI by Western blot at 0.5 to 2 ug/mL. Antibody can also be used for immunohistochemistry starting at 2 ug/mL. For immunofluorescence start at 20 ug/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Anti-OMI HTRA2 Antibody (A01941) Images



Western blot analysis of OMI in U937 lysate with Omi antibody at (A) 0.5, (B) 1, and (C) 2 ug/mL.



Immunohistochemistry of OMI in human liver tissue with OMI antibody at 2 ug/mL.



Immunofluorescence of OMI in Human Liver cells with OMI antibody at 20 ug/mL.

Anti-OMI HTRA2 Antibody

For Research Use Only. Not for use in diagnostic procedures.