

## Anti-UGT1A9 Antibody

Catalog Number: A01979

### About UGT1A9

Cullins assemble a potentially large number of ubiquitin ligases by binding to the RING protein ROC1 to catalyse polyubiquitination, as well as binding to various specificity factors to recruit substrates. Cullin 5 is a component of E3 ubiquitin ligase complexes, which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. Cullin 5 seems to be involved proteasomal degradation of p53/TP53 stimulated by adenovirus E1B-55 kDa protein. Cullin 5 may form a cell surface vasopressin receptor. Cullin 5 is part of a E3 ubiquitin ligase complex with elongin BC complex (TCEB1 and TCEB2), RBX1 and MUF1, complexes with elongin BC complex (TCEB1 and TCEB2), RXB1 and TCEB3 or SOCS1 or WSB1; elongin BC complex (TCEB1 and TCEB2), RBX1 and VHL; elongin BC complex (TCEB1 and TCEB2), RBX1, adenovirus type 5 E1B-55kDa protein and adenovirus type 5 E4-orf6. Cullin 5 Interacts with RBX1 and RNF7.

### Overview

Product Name	Anti-UGT1A9 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-UGT1A9 Antibody catalog # A01979. Tested in WB,IHC applications. This antibody reacts with Human,Mouse,Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O60656

### Technical Details

Immunogen	Synthesized peptide derived from human PC-PLD3
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

WB: 1:500-1:2000

IHC: 1:50-1:200

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-UGT1A9 Antibody