

Anti-Gastric Mucin/MUC6 Antibody Picoband®

Catalog Number: A02143

About MUC6

Mucin 6, oligomeric mucus/gel-forming, also known as MUC6, is a human gene. This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins produced by many epithelial tissues. The protein encoded by this gene is secreted and forms an insoluble mucous barrier that protects the gut lumen. The MUC6 gene encodes gastric mucin, a secreted glycoprotein that plays an essential role in epithelial cytoprotection from acid, proteases, pathogenic microorganisms, and mechanical trauma in the gastrointestinal tract.

Overview


Product Name	Anti-Gastric Mucin/MUC6 Antibody Picoband®
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Gastric Mucin/MUC6 Antibody Picoband® catalog # A02143. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q6W4X9

Technical Details


Immunogen	E. coli-derived human Gastric Mucin recombinant protein (Position: H52-R230).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml ELISA, 0.1-0.5ug/ml

Anti-Gastric Mucin/MUC6 Antibody Picoband® (A02143) Images



Western blot analysis of Gastric Mucin using anti-Gastric Mucin antibody (A02143). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: mouse stomach tissue lysates, Lane 2: human SGC-7901 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Gastric Mucin antigen affinity purified polyclonal antibody (Catalog # A02143) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Gastric Mucin at approximately 257KD. The expected band size for Gastric Mucin is at 257KD.



Western blot analysis of Gastric Mucin using anti-Gastric Mucin antibody (A02143). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: Normal group-rat colon tissue lysates, Lane 2: Model group-rat colon tissue lysates, Lane 3: Triditional Chinese medicine treatment (low dose)-rat colon tissue lysates, Lane 4: Triditional Chinese medicine treatment (medium dose)-rat colon tissue lysates, Lane 5: Triditional Chinese medicine treatment (high dose)-rat colon tissue lysates, Lane 6: Western medicine treatment-rat colon tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Gastric Mucin antigen affinity purified polyclonal antibody (Catalog # A02143) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with ChemiDoc MP system. A specific band was detected for Gastric Mucin at approximately 257KD. The expected band size for Gastric Mucin is at 257KD.

1 Publications Citing This Product

1. PubMed ID: 22027009, Mucins Differently Expressed in Various Ampullary Adenocarcinomas

Visit bosterbio.com/anti-gastric-mucin-antibody-a02143-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Gastric Mucin/MUC6 Antibody

For Research Use Only. Not for use in diagnostic procedures.