

## Anti-TAF1/Kat4 Antibody

Catalog Number: A02151

### About TAF1

Anti-TAF1 antibody is suitable for Cancer, Immunology and Nuclear Signaling research. Transcription Initiation Factor TFIID Subunit 1 (TAF1) is the largest component and core scaffold of the TFIID basal transcription factor complex, which also includes TATA-binding protein (TBP) and a variety of TBP-associated factors. TFIID nucleates the formation of transcription pre-initiation complexes and plays a key role in the regulation of gene expression by RNA polymerase II. TAF1 possesses DNA-binding activity and also contains novel N- and C-terminal Ser/Thr kinase domains which can auto-phosphorylate or trans-phosphorylate other transcription factors. For example, TAF1 interacts with the C-terminus of TP53 and phosphorylates the T55 residue, leading to MDM2-mediated degradation of TP53. TAF1 also catalyzes Ser phosphorylation of general transcription factor IIA (GTF2A1) and IIF (GTF2F1). The retinoblastoma tumor suppressor protein, RB1, interacts with the N-terminal domain of TAF1 and inhibits its intrinsic kinase activity. TAF1 is essential for progression through the G1 phase of the cell cycle and has been reported to be indispensable for expression of 18% of mammalian genes.

### Overview


Product Name	Anti-TAF1/Kat4 Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-TAF1/Kat4 Antibody (Catalog # A02151). Tested in CHIP, ELISA, WB applications. This antibody reacts with Human, Mouse.
Application	ChIP, ELISA, IP, IHC, WB
Clonality	Polyclonal
Formulation	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.01% (w/v) Sodium Azide
Storage Instructions	Store vial at -20°C prior to opening. Aliquot contents and freeze at -20°C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of opening. (Ship on dry ice.)
Host	Rabbit
Uniprot ID	P21675

### Technical Details

Immunogen	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to amino acids near the carboxyl terminus of human TAF1.
Predicted Reactive Species	Bovine, Pufferfish, Zebrafish
Isotype	IgG


Form	Liquid (sterile filtered)
Concentration	1.1 mg/mL by UV absorbance at 280 nm
Purification	This affinity purified antibody is directed against human TAF1 protein. The product was affinity purified from monospecific antiserum by immunoaffinity chromatography. A BLAST analysis was used to suggest cross-reactivity with TAF1 protein from human, mouse and rat based on 100% homology with the immunizing sequence. Reactivity against homologues from other sources is not known.
Suggested Dilutions	ELISA: 1:150,000 - 1:300,000 ChIP: User optimized IHC: 10µg/ml IP: User optimized WB: 1:100 - 1:2,000 This affinity purified antibody has been tested for use in ELISA, Immunohistochemistry, ChIP, and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 213 kDa in size corresponding to TAF1 protein by western blotting in the appropriate cell lysate or extract.

## Anti-TAF1/Kat4 Antibody (A02151) Images




Western Blot of Rabbit anti-TAF1 antibody. Marker: Opal Pre-stained ladder . Lane 1: HEK293 lysate . Lane 2: HeLa Lysate . Lane 3: MCF-7 Lysate . Lane 4: Jurkat Lysate . Lane 5: A431 Lysate . Lane 6: A549 Lysate . Lane 7: LNCap Lysate . Lane 8: MOLT-4 Lysate . Lane 9: Ramos Lysate . Lane 10: Raji Lysate . Lane 11: A-172 Lysate . Lane 12: NIH/3T3 Lysate . Load: 35  $\mu$ g per lane. Primary antibody: TAF1 antibody at 0.2 $\mu$ g/mL overnight at 4C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:30,000 for 60 min at RT. Blocking Buffer: 1% Casein-TTBS for 30 min at RT. Predicted/Observed size: 250kDa for TAF1.


Occupation of TAF1 on c-Fos Promoter after TNF Treatment



Transcription Initiation Factor TFIID Subunit 1 (TAF1) antibody was used to detect TAF1 in treated and untreated HeLa Cells. HeLa cells were treated with TNF alpha and Chromatin was prepared by EZ Magna Chip Kit (Millipore). CHIP was performed on fos promoters using 5  $\mu$ g of TAF1 antibody from Boster and an RNA PolII antibody. Image with data provided courtesy of Shiraz Mujtaba, Ph.D., Dept. of Structural & Chemical Biology, Mount Sinai School of Medicine.



Western blot using Boster's affinity purified anti-TAF1 to detect TAF1 in HeLa nuclear extract (arrowhead). The membrane was probed with the primary antibody at dilutions of 1:100 (lane 1), 1:250 (lane 2), 1: 500 (Lane 3 and 1:1,000 (Lane 4). The identity of the bands at ~95 kDa is unknown, but may be degraded TAF1. Personal Communication, Anne Gegonne, CCR-NCI, Bethesda, MD.



Immunohistochemistry of rabbit anti-TAF1 antibody. Tissue: prostate. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: Anti-TAF1 at 10  $\mu$ g/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Staining: TAF-1 as precipitated red signal with hematoxylin purple nuclear counterstain.

Western blot using Boster's affinity purified anti-TAF1 to detect TAF1 in HeLa nuclear extract (arrowhead). The membrane was probed with the primary antibody at dilutions of 1:100 (lane 1), 1:250 (lane 2), 1: 500 (Lane 3 and 1:1,000 (Lane 4). The identity of the bands at ~95 kDa is unknown, but may be degraded TAF1. Personal Communication, Anne Gegonne, CCR-NCI, Bethesda, MD.



## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-TAF1/Kat4 Antibody

For Research Use Only. Not for use in diagnostic procedures.