

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-NACA Antibody

Catalog Number: A02155

About NACA

Largest component and core scaffold of the TFIID basal transcription factor complex. Contains novel N- and C-terminal Ser/Thr kinase domains which can autophosphorylate or transphosphorylate other transcription factors. Phosphorylates TP53 on 'Thr-55' which leads to MDM2-mediated degradation of TP53. Phosphorylates GTF2A1 and GTF2F1 on Ser residues. Possesses DNA-binding activity. Essential for progression of the G1 phase of the cell cycle. Exhibits histone acetyltransferase activity towards histones H3 and H4.

Sekiguchi T., Mol. Cell. Biol. 11:3317-3325(1991). Ruppert S., Nature 362:175-179(1993). Nolte D., Proc. Natl. Acad. Sci. U.S.A. 100:10347-10352(2003).

Overview

Product Name	Anti-NACA Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-NACA Antibody catalog # A02155. Tested in WB,IHC applications. This antibody reacts with Human,Mouse,Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Rabbit IgG, $1mg/ml$ in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q13765

Technical Details

Immunogen	Synthesized peptide derived from internal of human TAF1.
Predicted Reactive Species	Boar, Bovine, Canine, Golden Hamster
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-



888-466-3604 | support@bosterbio.com | www.bosterbio.com

	specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:2000-1:5000 IHC: 1:50-1:200



888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-NACA Antibody (A02155) Images




Figure 1. Western blotting validation for Anti-NACA Antibody A02155

Western blot analysis of extracts of various cell lines

Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-NACA Antibody