

Anti-Human POR DyLight® 488 conjugated Antibody

Catalog Number: A02166-Dyl488

About POR

POR is a membrane-boundenzyme required for electron transfer from NADPH to cytochrome P450 in the endoplasmic reticulum of theeukaryotic cell. The gene encodes an endoplasmic reticulum membrane oxidoreductase with an FAD-binding domain and a flavodoxin-like domain. The protein binds two cofactors, FAD and FMN, which allow it to donate electrons directly from NADPH to all microsomal P450 enzymes. Mutations in this gene have been associated with various diseases, including apparent combined P450C17 and P450C21 deficiency, amenorrhea and disordered steroidogenesis, congenital adrenal hyperplasia and Antley-Bixler syndrome.

Overview

Product Name	Anti-Human POR DyLight® 488 conjugated Antibody
Reactive Species	Human
Description	Boster Bio Anti-Human POR DyLight® 488 conjugated Antibody catalog # A02166-Dyl488. Tested in Flow Cytometry applications. This antibody reacts with Human.
Conjugate	DyLight®488
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na2HPO4, 0.02% NaN3.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P16435

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human POR, different from the related mouse and rat sequences by five amino acids.
Predicted Reactive Species	Human
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Flow Cytometry, 1-3ug/1x10 ⁶ cells



Anti-Human POR DyLight® 488 conjugated Antibody (A02166-Dyl488) Images




Figure 1. Flow Cytometry analysis of A549 cells using anti-Human POR antibody (A02166-Dyl488). Overlay histogram showing A549 cells stained with A02166-Dyl488 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Human POR Antibody (A02166-Dyl488, 1ug/1x10⁶ cells) for 30 min at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Human POR DyLight® 488 conjugated Antibody